

# For Reference

---


**NOT TO BE TAKEN FROM THIS ROOM**

# For Reference

---

NOT TO BE TAKEN FROM THIS ROOM

EX LIBRIS  
UNIVERSITATIS  
ALBERTAENSIS











SOME OPERATOR NORMALITIES ON BANACH SPACES

by

TWUM OWUSU-ANSAH



A THESIS

SUBMITTED TO THE FACULTY OF GRADUATE STUDIES  
IN PARTIAL FULFILMENT OF THE REQUIREMENTS FOR THE DEGREE  
OF MASTER OF SCIENCE

DEPARTMENT OF MATHEMATICS

UNIVERSITY OF ALBERTA

EDMONTON, ALBERTA

MAY, 1968





UNIVERSITY OF ALBERTA  
FACULTY OF GRADUATE STUDIES

The undersigned certify that they have  
read and recommend to the Faculty of Graduate Studies  
for acceptance, a thesis entitled "SOME OPERATOR  
NORMALITIES ON BANACH SPACES", submitted by  
TWUM OWUSU-ANSAH in partial fulfilment of the  
requirements for the degree of Master of Science.



(i)

ABSTRACT

The introduction of semi-inner product spaces made possible two definitions of normal operators on a Banach space. In this paper we find the relation between the two kinds of normal operators and investigate some of their simple properties. We also use the semi-inner product in an attempt to extend the concept of hyponormal operators to Banach spaces.



ACKNOWLEDGEMENTS

The author is greatly indebted to Professor J. S. W. Wong for suggesting the topic that led to this thesis and for his helpful hints, criticisms and sustained interest in its preparation; to Professors D. W. Boyd and J. W. Macki who in the absence of Professor Wong provided useful criticism and encouragement; and to the Canadian Commonwealth Scholarship and Fellowship Committee for giving him the opportunity and assistance that made the preparation of this thesis possible.



TABLE OF CONTENTS

	Page
ABSTRACT . . . . .	(i)
ACKNOWLEDGEMENTS . . . . .	(ii)
GLOSSARY . . . . .	(iv)
 CHAPTER I:      PRELIMINARY RESULTS AND NOTATION . . . . .	 1
 CHAPTER II:     PSEUDONORMAL AND NORMAL OPERATORS . . . . .	 17
 CHAPTER III:    HYPONORMAL OPERATORS . . . . .	 40
 BIBLIOGRAPHY . . . . .	 73





(iv)

## GLOSSARY

<u>SYMBOL OR NAME</u>	<u>DEFINITION</u>
$f(\alpha) = o(\alpha)$ as $\alpha \rightarrow 0$ ,	$\frac{ f(\alpha) }{\alpha} \rightarrow 0$ as $\alpha \rightarrow 0$
$\mathcal{C}(A)$ , convex hull of a set $A$ ,	The smallest convex set containing $A$ .
$\rho(T)$ , resolvent set of an operator $T$ on a space $X$ .	The set of all complex numbers $\lambda$ such that $(\lambda I - T)^{-1}$ exists as a bounded linear operator with domain $X$ .
$\sigma(T)$ , the spectrum of $T$ ,	The complement in the complex plane of the resolvent set of $T$ .
$P\sigma(T)$ , the point spectrum of $T$ ,	The subset of $\sigma(T)$ for which $(\lambda I - T)^{-1}$ is not one-to-one.
$r(T)$ , the spectral radius of $T$ ,	$\sup \{  \lambda  : \lambda \in \sigma(T) \}$ .
$M_1 \oplus M_2$ , the direct sum of the subspaces $M_1, M_2$ of a space $X$	The smallest subspace in $X$ containing both $M_1$ and $M_2$ where $M_1 \cap M_2 = \{0\}$ .
$\sigma$ -Algebra of sets,	A family $F$ of subsets of a set $X$ such that $A_k, A, B \in F \implies A \cap B \in F$ , $X - A \in F$ and $\bigcup_{k=1}^{\infty} A_k \in F$ .
Borel $\sigma$ -Algebra of plane sets,	The smallest $\sigma$ -Algebra of subsets of the complex plane containing all rectangles.



<u>SYMBOL OF NAME</u>	<u>DEFINITION</u>
$T^*$ , the Hilbert space adjoint of an operator $T$ ,	$(Tx, y) = (x, T^*y)$
Projection	A bounded linear operator on a Banach space $X$ such that $P^2 = P$ . In the case of a Hilbert space the condition $P^* = P$ is also required.
Algebra	A ring $R$ which is a linear space and satisfies the condition $(\alpha x)y = \alpha(xy) = x(\alpha y)$ for $\alpha \in \mathbb{C}$ , $x, y \in R$ .
Division Algebra	An algebra such that every element $T \neq 0$ , has an inverse.
Boolean Algebra of projections	A distributive lattice $R$ of projections containing two elements $0$ and $I$ such that (i) $0 \leq x \leq I$ for all $x \in R$ (ii) for any $x \in R$ , there is some $x' \in R$ such that $x \vee x' = I$ , $x \wedge x' = 0$ .
Nilpotent operator $T$ ,	An operator $T$ for which there is some integer $n$ such that $T^n = 0$ .
Generalized quasi-nilpotent operator $T$	An operator $T$ such that $\lim_{n \rightarrow \infty} \ T^n\ ^{1/n} = 0$ .
A positive operator $T$ in a Hilbert space $X$ , $T \geq 0$ ,	An operator $T$ such that $(Tx, x) \geq 0$ for all $x \in X$ .



## CHAPTER I

### PRELIMINARY RESULTS AND NOTATION

1.1  $X$  will denote a complex Banach space and  $B(X)$  will denote the associated Banach algebra of bounded linear operators on  $X$ .

#### 1.2 Semi-inner product spaces

A semi-inner product on a linear space  $Y$  is a mapping  $[\cdot, \cdot]$  from  $Y \times Y$  into  $\mathbb{C}$ , the complex field such that

$$\begin{aligned} 1.2.1 \quad & [x+y, z] = [x, z] + [y, z] \\ & [\lambda x, y] = \lambda [x, y], \text{ for all complex } \lambda \text{ and} \\ & \text{all } x, y, z \in Y. \end{aligned}$$

$$1.2.2 \quad [x, x] > 0 \text{ if } x \neq 0, \quad x \in Y$$

$$1.2.3 \quad |[x, y]|^2 \leq [x, x][y, y], \quad x \text{ and } y \in Y.$$

A semi-inner product space (S.I.P.S) is a linear space on which a semi-inner product is defined. These definitions were introduced by G. Lumer [12] who proved the following results

$$\begin{aligned} 1.2.4 \quad & \text{Every semi-sinner product space is normed, the norm of} \\ & \text{a point } x \text{ being } [x, x]^{1/2} \end{aligned}$$



1.2.5 Every normed linear space can be made into a semi-inner product space in such a way that the norm induced by the semi-inner product on the normed space coincides with the norm on the space. [In general this can be done in infinitely many ways. There is a unique semi-inner product on a normed space if and only if the unit sphere of the space is smooth. In particular the unique semi-inner product on a Hilbert space is the inner product.]

1.3 The numerical range of an operator  $T \in B(X)$  .

The set of complex numbers

$$W(T) = \{[Tx, x] : [x, x] = 1, x \in X\} .$$

where  $[\cdot, \cdot]$  denotes one of the semi-inner products that can be introduced on the space  $X$  in the manner of 1.2.5, is called a numerical range for the operator  $T$  . It follows that there are many numerical ranges of an operator in general, depending on which semi-inner product is introduced on  $X$  . This definition is due to G. Lumer [12] who also proved the following results:

1.3.1 The convex hull of a numerical range for an operator on a normed linear space is independent of the semi-inner product used, as long as the semi-inner product yields the norm.

1.3.2 In particular, if the numerical range of an operator  $T$  is real for one semi-inner product introduced on  $X$  in the





manner of 1.2.5, then it is real for all other semi-inner products.

#### 1.4 Hermitian Operators

An operator  $T \in B(X)$  is said to be hermitian if and only if its numerical range with respect to some semi-inner product introduced on  $X$  in the manner of 1.2.5 is real. By virtue of 1.3.2, it is clear that this concept of hermiticity does not depend on the semi-inner product used. The following result is due to G. Lumer [12].

1.4.1 An operator  $T \in B(X)$  is hermitian if and only if

$$\|I + i\alpha T\| = 1 + o(\alpha) \text{ as } \alpha \rightarrow 0,$$

where  $\alpha$  is real and  $I$  is the identity operator in  $B(X)$ .

This result establishes the equivalence of G. Lumer's definition of hermiticity and that given by I. Vidav [19].

#### 1.5 Reducibility of an operator $T \in B(X)$

An operator  $T \in B(X)$  is said to be completely reduced by a pair of sub-spaces  $M_1, M_2$  of  $X$  if and only if

$$1.5.1 \quad X = M_1 \oplus M_2$$



$$1.5.2 \quad TM_1 \subset M_1, \quad TM_2 \subset M_2.$$

(See [18, page 268].)

## 1.6 Decomposition Theorem.

Let  $T \in B(X)$ . An admissible domain for  $T$  is any bounded open set  $D$  in the complex plane whose boundary  $\partial D$  consists of a finite number of closed rectifiable curves lying in the resolvent set  $\rho(T)$  of  $T$  and oriented according to the usual convention in the complex plane. An isolated part  $\sigma$  of  $\sigma(T)$ , the spectrum of  $T$ , is any subset of  $\sigma(T)$  which is at a positive distance from its complementary part  $\sigma' = \sigma(T) - \sigma$ . If  $\sigma$  is an isolated part of  $\sigma(T)$ , then there exists an admissible domain  $D$  for  $T$  such that  $\sigma = \sigma(T) \cap D$ . In this case we have the following decomposition theorem [16, p. 420-421].

There exist subspaces  $M_\sigma, M_{\sigma'}$  of  $X$  such that

$$1.6.1 \quad X = M_\sigma \oplus M_{\sigma'}$$

$$1.6.2 \quad TM_\sigma \subset M_\sigma, \quad TM_{\sigma'} \subset M_{\sigma'}$$

$$1.6.3 \quad \text{Spectrum of } T|_{M_\sigma} = \sigma$$

$$1.6.4 \quad \text{Spectrum of } T|_{M_{\sigma'}} = \sigma(T) - \sigma = \sigma'$$



### 1.7 Spectro-normaloid operators

An operator  $T \in B(X)$  such that  $\|T\| = \max \{ |\lambda| : \lambda \in \sigma(T) \}$ , where  $\sigma(T)$  denotes the spectrum of  $T$ , will be called spectro-normaloid. In Hilbert space an operator  $T$  is spectro-normaloid if and only if it is normaloid [10, p. 416], i.e.  $\|T\| = \max \{ |\lambda| : \lambda \in W(T) \}$ .

### 1.8 The property SN

A set of operators  $S \subset B(X)$  such that every member is spectro-normaloid will be said to have property SN.

### 1.9 Spectral Operators of scalar type

Let  $\mathcal{A}$  be the  $\sigma$ -algebra of Borel subsets of the complex plane  $\mathbb{C}$ . Let  $X$  be an arbitrary Banach space over the complex field. The function  $p(\cdot) : \mathcal{A} \rightarrow B(X)$  is called a spectral measure in  $X$  iff the following hold for all  $\Delta_1, \Delta_2, \Delta_3 \in \mathcal{A}$ :

$$1.9.1 \quad p(\Delta_1 \cap \Delta_2) = p(\Delta_1)p(\Delta_2)$$

$$1.9.2 \quad p(\Delta_1 \cup \Delta_2) = p(\Delta_1) + p(\Delta_2) - p(\Delta_1)p(\Delta_2)$$

$$1.9.3 \quad p(\mathbb{C} - \Delta) = I - p(\Delta), \text{ where } I \text{ is the identity operator in } B(X).$$

$$1.9.4 \quad \text{There exists a constant } M \text{ such that } \|p(\Delta)\| \leq M \text{ for all } \Delta \in \mathcal{A}.$$



It follows from 1.9.1 that  $p(\Delta) = \{p(\Delta)\}^2$  for all  $\Delta \in \mathcal{A}$  and hence that  $p(\cdot)$  is projection-valued. Thus the range of  $p(\cdot)$  is a bounded Boolean algebra of projections.

A subset  $\Gamma$  of  $X'$ , the conjugate space of  $X$ , is said to be total if  $f(x) = 0$  for every  $f \in \Gamma$  implies  $x = 0$ . If  $\Gamma$  is a total subset of  $X'$ , then the spectral measure  $p(\cdot)$  is said to be  $\Gamma$ -completely additive if

1.9.5 For any pairwise disjoint sequence  $\{\Delta_k\}_{k=1}^{\infty}$  of  $\mathcal{A}$  we have for all  $x \in X$  and all  $f \in \Gamma$ ,

$$f\{p(\bigcup_{k=1}^{\infty} \Delta_k)x\} = \sum_{k=1}^{\infty} f\{p(\Delta_k)x\}$$

An operator  $T \in B(X)$  is called a spectral operator if there exists a  $\Gamma$ -completely additive spectral measure  $p(\cdot)$  in  $X$  such that the following conditions hold:

1.9.6  $T$  commutes with  $p(\Delta)$  for all  $\Delta \in \mathcal{A}$ .

1.9.7 Let  $X_{\Delta} = p(\Delta)X$  and  $T_{\Delta} = p(\Delta)T$ . If  $u \in X_{\Delta}$ , then  $u = p(\Delta)x$  for some  $x \in X$ . Hence

$$\begin{aligned} T_{\Delta}u &= p(\Delta)Tu \\ &= p(\Delta)Tp(\Delta)x \\ &= p(\Delta)^2Tx, \end{aligned}$$

assuming 1.9.6 holds. That is  $T_{\Delta}u = p(\Delta)Tx$  since





$$\{p(\Delta)\}^2 = p(\Delta) .$$

Thus  $T_{\Delta}u = p(\Delta)(Tx) \in X_{\Delta}$  . Hence  $T_{\Delta}u \in X_{\Delta}$  and so  $T_{\Delta}$  maps  $X_{\Delta}$  into itself. That is  $T_{\Delta} \in B(X_{\Delta})$ . The condition to be satisfied is that the spectrum of  $T_{\Delta}$  considered as an element of  $B(X_{\Delta})$  must be contained in the closure of  $\Delta$  .

When  $T$  is a spectral operator the spectral measure  $p(\cdot)$  is called a resolution of the identity for  $T$  . For such an operator  $T$  , N. Dunford [5, pages 329-330] has proved the following results:

1.9.8  $p(\cdot)$  is unique

1.9.9 Any operator which commutes with  $T$  , commutes with  $p(\Delta)$  for all  $\Delta \in \mathcal{A}$  .

Let  $T$  be a spectral operator with resolution of the identity  $p(\cdot)$  . Let  $\sigma(T)$  , the spectrum of  $T$  , which is compact, be partitioned into disjoint non-void Borel sets  $\Delta_1, \Delta_2, \dots, \Delta_n$  whose diameters are each less than  $\delta$  ,  $\delta$  being an arbitrary positive number. Let us define  $\|\pi\|$  , the norm of such a partition  $\pi = (\Delta_1, \Delta_2, \dots, \Delta_n)$  as  $\max_{1 \leq i \leq n} \{\text{diam } \Delta_i\}$  . For a scalar-valued or operator-valued function  $f$  defined on  $\sigma(T)$  if

$$\lim_{\|\pi\| \rightarrow 0} \sum_{i=1}^n f(\lambda_i) p(\Delta_i)$$



exists independently of the choice of  $\pi$  and  $\lambda_i \in \Delta_i$   
 $\{i = 1, 2, \dots, n\}$ , we say that  $f$  is integrable and define

$$\int_{\sigma(T)} f(\lambda) p(d\lambda) ,$$

the integral of  $f$ , as the limit

$$\lim_{\|\pi\| \rightarrow 0} \sum_{i=1}^n f(\lambda_i) p(\Delta_i)$$

which is taken in the uniform operator topology. Thus

$$1.9.10 \quad \int_{\sigma(T)} f(\lambda) p(d\lambda) = \lim_{\|\pi\| \rightarrow 0} \sum_{i=1}^n f(\lambda_i) p(\Delta_i)$$

A spectral operator  $T$  such that

$$T = \int_{\sigma(T)} \lambda p(d\lambda)$$

is said to be of scalar type.

## 1.10 Gelfand Theory of Commutative Banach Algebras

A Banach algebra  $Y$  is an algebra which is also Banach space such that  $\|x y\| \leq \|x\| \|y\|$  for all  $x, y \in Y$ . Let  $Y$  denote a complex commutative Banach algebra with identity  $e$  of norm 1. A non-empty subset  $I$  of  $Y$  is called an ideal if

1.10.1  $I$  is a subspace of  $Y$



$$1.10.2 \quad x \in I, y \in Y \implies xy \in I$$

An ideal  $I$  is proper if  $I \neq Y$ . A proper ideal  $M$  is said to be maximal if the only ideal that contains  $M$  properly is all of  $Y$ . A maximal ideal is closed. Since  $Y$  is a complex commutative Banach algebra with identity  $e$ , it follows that the quotient algebra  $Y/M$  is also a complex commutative Banach algebra with identity  $e + M$ ,  $M$  being any maximal ideal.

Now  $Y$  is also a commutative ring with identity and since  $M$  is a maximal ideal in  $Y$ , it follows by the Gelfand-Mazur theorem that since  $Y/M$  is a complex Banach algebra with identity as well as a division algebra, it is isomorphic to the complex field. It is also clear from the proof of this result [1, p. 42] that given  $y$  and  $M$ , there exists a complex number  $y(M)$  which corresponds to  $y + M$  in the isomorphism, such that  $y + M = y(M) (e + M)$ . Thus using the natural homomorphism  $Y \rightarrow Y/M$  we may map  $Y$  onto the complex field as follows

$$Y \rightarrow Y/M \rightarrow \text{complex field}$$

$$\text{where } y \rightarrow y + M \rightarrow y(M).$$

Suppose we now fix  $y$  and let  $M$  vary over the set of all maximal ideals  $\mathcal{M}$  in  $Y$ . Then  $y$  determines a unique function  $\hat{y}$  defined over  $\mathcal{M}$  such that  $\hat{y}(M) = y(M)$ . Thus if we let  $C(\mathcal{M})$  denote the set of all these complex-valued functions  $\hat{y}$ ,



we obtain a homomorphism from  $Y$  into the algebra of complex-valued functions  $C(\mathcal{M})$ .

This homomorphism has the following properties:

$$1.10.3 \quad \widehat{x_1 + x_2} = \widehat{x_1} + \widehat{x_2}$$

$$1.10.4 \quad \widehat{\alpha x} = \alpha \widehat{x}$$

$$1.10.5 \quad \widehat{x_1 x_2} = \widehat{x_1} \widehat{x_2}$$

$$1.10.6 \quad \widehat{e} = 1$$

We also have the properties

$$1.10.7 \quad \sigma(y) = \{y(M) : M \in \mathcal{M}\}$$

$$1.10.8 \quad r(y) = \sup \{|y(M)| : M \in \mathcal{M}\}$$

The set  $\mathcal{M}$  can be topologised by giving it the weakest topology with respect to which all the functions  $\{\widehat{y}\}_{y \in Y}$  are continuous. This is the Gelfand topology on  $\mathcal{M}$ . With respect to this topology we have the following results:

$$1.10.9 \quad \mathcal{M} \text{ is a compact Hausdorff space}$$

$$1.10.10 \quad \text{All the functions } \{\widehat{y}\}_{y \in Y} \text{ are continuous on } \mathcal{M}.$$





1.11 Some results due to E. Berkson

1.11.1 If  $A$  is a commutative Banach algebra and if every  $a \in A$  can be written in the form  $a = s + it$ , where  $s$  and  $t$  are hermitian operators belonging to  $A$ , then for all  $a \in A$ , we have  $\|a\| = r(a)$ , the spectral radius of  $a$ . [3, p. 5]

1.11.2 Suppose  $X$  is reflexive. If  $T$  is a spectral operator of scalar type then there exist operators  $u, v \in B(X)$  such that (i)  $T = u + iv$  (ii)  $uv = vu$ , (iii) relative to some norm on  $X$  equivalent to the given norm,  $u^m v^n$  are all hermitian for  $m, n = 0, 1, 2, \dots$ .

[2, p. 371]

1.11.3 If  $X$  is reflexive and  $T = u + iv$  where  $u, v \in B(X)$ ,  $uv = vu$  and relative to some norm on  $X$  equivalent to the given one, we have  $u^m v^n$  hermitian for  $m, n = 0, 1, 2, \dots$ , then  $T$  is spectral of scalar type.

[2, p. 371]

1.12 A result due to S. R. Foguel

If  $T \in B(X)$  is spectral of scalar type, then there exist operators  $R, S \in B(X)$  such that

(i)  $T = RS = SR$

(ii)  $\sigma(R)$  is a set of non-negative real numbers and  $\sigma(S)$  is a



subset of the unit circle.

[Theorem 2, 7, p. 61]

### 1.13 Normal Operators on a Hilbert Space

If  $T$  is a normal operator on a Hilbert space, the following hold (see [16] and [18]):

1.13.1  $T = \int_{\sigma(T)} \lambda p(d\lambda)$  where  $p(\cdot)$  is the resolution of the identity for  $T$ .

1.13.2  $\lambda T$  and  $\lambda + T$  are normal for all complex  $\lambda$

1.13.3  $\|T\| = r(T)$

1.13.4  $T^n$  is normal for all non-negative integral  $n$

1.13.5 There exist operators  $R, S$  where  $R$  is unitary and  $S$  is positive such that

$$T = RS = SR$$

1.13.6  $T^*$ , the adjoint of  $T$  is normal and conversely.

1.13.7 Any operator which commutes with  $T$ , commutes with  $p(\Delta)$  for all  $\Delta$ .

1.13.8 If  $S$  is an operator such that  $ST = TS$ , then

}  $S + T$  and  $TS$  are both normal.



1.13.9 If a subspace  $M$  of  $X$  reduces  $T$ , then  $T|_M$  is normal.

1.13.10 The limit in the uniform operator topology of a sequence of normal operators is normal.

#### 1.14 Hyponormal operators on a Hilbert space .

An operator  $T$  on a Hilbert space  $H$  is said to be hyponormal if  $\|T^*x\| \leq \|Tx\|$  for all  $x \in H$ . The following results hold:

1.14.1  $T$  is hyponormal if and only if

$$T^*T - TT^* \geq 0 .$$

Proof:

Suppose  $T$  is hyponormal. Then for all  $x \in H$ ,

$$\begin{aligned} (T^*Tx, x) &= (Tx, Tx) = \|Tx\|^2 \geq \|T^*x\|^2 \\ &= (T^*x, T^*x) \\ &= (TT^*x, x) \end{aligned}$$

$$\text{i.e. } T^*T - TT^* \geq 0 .$$

Now suppose  $T^*T - TT^* \geq 0$ . Then for all  $x \in H$ ,

$$\begin{aligned} \|T^*x\|^2 &= (T^*x, T^*x) = (TT^*x, x) \leq (T^*Tx, x) \\ &= (Tx, Tx) \\ &= \|Tx\|^2 \end{aligned}$$

i.e.  $T$  is hyponormal.



1.14.2 If  $T$  is hyponormal then so are  $\lambda T$  and  $T + \lambda$  for all complex  $\lambda$ .

Proof:

Since  $T$  is hyponormal, we have

$$([T^*T - TT^*]x, x) \geq 0 \quad \text{for all } x \in X.$$

Now

$$\begin{aligned} & ([(\lambda T)^*(\lambda T) - (\lambda T)(\lambda T)^*]x, x) \\ &= ([\bar{\lambda} T^* \lambda T - \lambda T \bar{\lambda} T^*]x, x) \\ &= |\lambda|^2 ([T^*T - TT^*]x, x) \geq 0. \end{aligned}$$

Hence  $\lambda T$  is hyponormal

$$\begin{aligned} & ([T+\lambda]^*(T+\lambda) - (T+\lambda)(T+\lambda)^*)x, x) \\ &= ([T^* + \bar{\lambda}](T+\lambda) - (T+\lambda)(T^* + \bar{\lambda}'))x, x) \\ &= ([T^*T - TT^*]x, x) \geq 0. \end{aligned}$$

Hence  $T + \lambda$  is hyponormal.

1.14.3 If  $T$  is hyponormal, then  $\|T\| = r(T)$ , the spectral radius of  $T$ ,

[Theorem 1, p. 1453, 17]

1.14.4 If  $T$  is hyponormal and  $M \subset X$  is invariant under  $T$ , then  $T|_M$  is also hyponormal.





Proof:

$$\|T^*x\| \leq \|Tx\| \quad \text{for all } x \in H .$$

In particular,

$$\|T^*x\| \leq \|Tx\| \quad \text{for all } x \in M .$$

Hence we have

$$1.14.4.1 \quad \|T^*|_M\| \leq \|T|_M\| .$$

Now  $M \subset X$  is closed subspace and so is a Hilbert space. It follows then that

$$(T|_M)^* : M^* \rightarrow M^*$$

and so

$$(T|_M)^* : M \rightarrow M .$$

But

$$T^* : X^* \rightarrow X^* ,$$

that is

$$T^* : X \rightarrow X .$$

Thus

$$T^*|_M : M \rightarrow M .$$

Let  $x \in M$  be arbitrary. Then for all  $y \in M$  we have

$$\begin{aligned} ([T|_M]^*x, y) &= (x, [T|_M]y) \\ &= (x, Ty) \\ &= (T^*x, y) \\ &= (T^*|_Mx, y) . \end{aligned}$$



In particular, putting  $y = (T|_M)^* x \in M$  we have

$$([T|_M]^* x, [T|_M]^* x) = (T^*|_M x, [T|_M]^* x) \quad .$$

Thus

$$\begin{aligned} \|(T|_M)^* x\|^2 &= |(T^*|_M x, [T|_M]^* x)| \\ &\leq \|T^*|_M x\| \cdot \|(T|_M)^* x\| \end{aligned}$$

by Schwarz's inequality

$$\leq \|(T|_M)x\| \cdot \|(T|_M)^* x\| \quad ,$$

by 1.14.4.1.

Hence

$$\|(T|_M)^* x\| \leq \|(T|_M)x\|$$

for all  $x \in M$  and so  $T|_M$  is hyponormal.



## CHAPTER II

### PSEUDONORMAL AND NORMAL OPERATORS

#### 2.1 Pseudonormal Operators

An operator  $T \in B(X)$  will be called pseudonormal if and only if it can be written in the form  $T = u + iv$  where  $u$  and  $v$  are hermitian operators in  $B(X)$  such that  $uv = vu$ . Lumer [13] called these operators normal and showed that  $u$  and  $v$  are unique. We shall call  $u$  and  $v$  the components of the pseudonormal operator  $T$ .

#### 2.2 Normal Operators

An operator  $T \in B(X)$  will be called normal if and only if it can be written in the form

$$T = u + iv, \quad \text{where} \quad uv = vu,$$

$u$  and  $v \in B(X)$  and  $u^m v^n$  are all hermitian operators for  $m, n = 0, 1, 2, \dots$ . This definition is due to E. Berkson [3]. Clearly normal operators are pseudonormal.



2.3 Theorem: Let  $u, v \in B(X)$  be two commuting operators. Let  $B_o(X)$  be the closed subalgebra of  $B(X)$  generated by  $I, u$  and  $v$ . Then  $u$  and  $v$  hermitian implies  $u^m v^n$  hermitian for all  $m, n = 0, 1, 2, \dots$  if and only if  $B_o(X)$  has property SN.

Proof:

(i) Sufficiency

$B_o(X)$  is a commutative Banach algebra with identity. Hence by the Gelfand representation theory (1.10) we have a homomorphism from  $B_o(X)$  into  $C(\mathcal{M})$ , the space of continuous functions over the maximal ideal space  $\mathcal{M}$ . Since  $u$  and  $v$  are hermitian operators,  $\sigma(u)$  and  $\sigma(v)$  are real [9, p. 548]\*. Thus by 1.10.7 we have

2.3.1  $U(M)$  and  $V(M)$  are real for all  $M \in \mathcal{M}$

Let  $t$  be real. Since  $I + it u^m v^n \in B_o(X)$  and  $B_o(X)$  has property SN, it follows that  $I + it u^m v^n$  is spectro-normaloid and so by 1.10.8 we have

$$\begin{aligned} \|I + it u^m v^n\| &= r\{I + it u^m v^n\} \\ &= \sup_{M \in \mathcal{M}} |(I + it u^m v^n)(M)|. \end{aligned}$$

---

\* This also follows from a more general result of J. P. Williams. See [20],





Thus

$$2.3.2 \quad \|I + it u^m v^n\| = \sup_{M \in \mathcal{M}} |(I + it u^m v^n)(M)|$$

Since  $(I + it u^m v^n)(M)$  is a complex-valued continuous function over the compact Hausdorff space  $\mathcal{M}$ , by 1.10.9 and 1.10.10, it follows that  $|(I + it u^m v^n)(M)|$  is a real-valued continuous function over  $\mathcal{M}$ . Hence there exists  $M_0 \in \mathcal{M}$  such that

$$\sup_{M \in \mathcal{M}} |(I + it u^m v^n)(M)| = |(I + it u^m v^n)(M_0)|$$

Hence by 1.10.3 to 1.10.6 we have

$$\begin{aligned} \sup_{M \in \mathcal{M}} |(I + it u^m v^n)(M)| &= |(I + it u^m v^n)(M_0)| \\ &= |I(M_0) + it \{u(M_0)\}^m \{v(M_0)\}^n| \\ &= |1 + it \lambda|, \end{aligned}$$

where  $\lambda = \{u(M_0)\}^m \{v(M_0)\}^n$  is real by 2.3.1. Hence by 2.3.2 we have

$$\begin{aligned} \|I + it u^m v^n\| &= |1 + it \lambda| \\ &= \sqrt{1 + t^2 \lambda^2}, \quad \text{since } \lambda t \text{ is real} \\ &= 1 + \frac{1}{2} t^2 \lambda^2 + \dots \\ &= 1 + o(t), \quad \text{as } t \rightarrow 0. \end{aligned}$$



Thus by the equivalence of Lumer's and Vidav's hermiticity [1.4.1] it follows that  $u^m v^n$  are hermitian operators for all  $m, n = 0, 1, 2, \dots$ .

(ii) Necessity

We shall use the following lemma which was essentially established by E. Berkson [2]. A variant of his proof will be given here for the sake of completeness:

2.3.3 Lemma

If  $B$  is a subset of  $B(X)$  such that every  $b \in B$  can be written in the form  $b = s + it$  where  $s$  and  $t$  are hermitian operators in  $B$ , then every  $c \in \overline{B}$ , the closure of  $B$ , can be written in the form  $c = s_0 + it_0$  where  $s_0$  and  $t_0$  are hermitian operators in  $\overline{B}$ .

By hypothesis, we have  $uv = vu$  and  $u^m v^n$  hermitian for all  $m, n = 0, 1, 2, \dots$ .

Let  $B$  denote the class of polynomials in  $u$  and  $v$  with complex coefficients. Then  $B$  is a commutative subalgebra of  $B(X)$ .  $\overline{B}$ , being the closed subalgebra of  $B(X)$  generated by  $I$ ,  $u$  and  $v$  is also commutative. Thus  $\overline{B} = B_0(X)$  is a commutative Banach algebra. Since every  $P \in B$  is a polynomial in  $u$  and  $v$  with complex coefficients, each  $P \in B$  can be written  $P = R + iJ$ , where  $R$  and  $J$  are polynomials in  $u$  and  $v$  with real coefficients. Since



$u^n v^m$  are all hermitian for  $m, n = 0, 1, 2, \dots$ ,  $R$  and  $J$  are also hermitian. Furthermore  $R$  and  $J$  belong to  $B$ .

Hence by lemma 2.3.3, every element  $C \in \overline{B} = B_0(X)$  can be written in the form  $C = R_0 + i J_0$  where  $R_0$  and  $J_0$  are hermitian and belong to  $B_0(X)$ . Thus  $B_0(X)$  is a commutative Banach algebra such that every  $C \in B_0(X)$  can be written in the form  $C = R_0 + i J_0$  where  $R_0$  and  $J_0$  are hermitian operators in  $B_0(X)$ . It follows by 1.11.1 that for all  $T \in B_0(X)$  we have

$$\|T\| = r(T) \quad .$$

Thus  $B_0(X)$  has property SN.

Now to prove lemma 2.3.3, let  $C \in \overline{B}$ . Then there exists a sequence  $\{c_n\}_{n=1}^{\infty} \in B$  such that  $c_n \rightarrow c$  as  $n \rightarrow \infty$ .

Since  $c_n \in B$ , it follows that  $c_n = s_n + i t_n$  where  $s_n$  and  $t_n$  are hermitian operators belonging to  $B$ .

Let  $\|x\| = 1 = [x, x]^{1/2}$

Then

$$\begin{aligned} & |[(s_n - s_m)x, x] + i[t_n - t_m]x, x]| \\ &= |[(s_n + it_n) - (s_m + it_m)]x, x]| \\ &= |[(c_n - c_m)x, x]| \quad . \end{aligned}$$

Hence by 1.2.3, we have



$$\begin{aligned} 2.3.5 \quad & |[(s_n - s_m)x, x] + i[(t_n - t_m)x, x]| \\ & \leq \|c_n - c_m\| . \end{aligned}$$

Since  $s_n, s_m, t_n, t_m$  are all hermitian, so are  $s_n - s_m$  and  $t_n - t_m$ .  
Hence  $[(s_n - s_m)x, x], [(t_n - t_m)x, x]$  are both real. Thus

$$\begin{aligned} & |[(s_n - s_m)x, x] + i[(t_n - t_m)x, x]| \\ & = \sqrt{[(s_n - s_m)x, x]^2 + [(t_n - t_m)x, x]^2} . \end{aligned}$$

It follows by 2.3.5 that we have

$$\begin{aligned} 2.3.6 \quad & |[(s_n - s_m)x, x]| \leq \|c_n - c_m\| \\ & |[(t_n - t_m)x, x]| \leq \|c_n - c_m\| \end{aligned}$$

But for all operators  $T \in B(X)$  we have [2, p. 366]

$$\sup_{\|x\|=1} |[Tx, x]| \leq \|T\| \leq 4 \sup_{\|x\|=1} |[Tx, x]| .$$

Thus by 2.3.6 we have

$$\begin{aligned} 2.3.7 \quad & \|s_n - s_m\| \leq 4\|c_n - c_m\| \\ & \|t_n - t_m\| \leq 4\|c_n - c_m\| . \end{aligned}$$





Since  $\{c_n\}_{n=1}^{\infty}$  is convergent, it follows by 2.3.7 that both sequences  $\{s_n\}_{n=1}^{\infty}$ ,  $\{t_n\}_{n=1}^{\infty}$  are convergent.

Let  $\lim_{n \rightarrow \infty} t_n = t_0$ ,  $\lim_{n \rightarrow \infty} s_n = s_0$ .

Then  $c = \lim_{n \rightarrow \infty} c_n = s_0 + it_0$ .

Let  $\epsilon$  be an arbitrary positive real number. Then there exists  $N = N(\epsilon)$ , a positive integer, such that for all  $m > N$  we have

$$2.3.8 \quad \|s_0 - s_m\| < \epsilon, \quad \|t_0 - t_m\| < \epsilon.$$

$$\begin{aligned} \text{Hence } \|I + its_0\| &= \|I + its_m - its_m + its_0\| \\ &\leq \|I + its_m\| + |t| \|s_0 - s_m\| \\ &\leq 1 + o(t) + |t| \epsilon \text{ as } t \rightarrow 0. \end{aligned}$$

Since  $\epsilon$  is arbitrary and the left hand side is independent of it, it follows that

$$\|I + its_0\| \leq 1 + o(t) \text{ as } t \rightarrow 0.$$

By a remark in [19, p. 122] this implies

$$\|I + its_0\| = 1 + o(t) \text{ as } t \rightarrow 0.$$

Hence  $s_0$  is a hermitian operator. Similarly  $t_0$  is a hermitian operator.



Since  $t_n$  and  $s_n$  belong to  $B$ ,  $s_0$  and  $t_0$  belong to  $\overline{B}$ . Thus  $c = s_0 + it_0$  where  $s_0$  and  $t_0$  are hermitian operators belonging to  $\overline{B}$ .

## 2.4 Remarks

2.4.1 If  $X$  is a Hilbert space, then since the product of any two commuting hermitian operators is hermitian, we have by 2.3 that the closed subalgebra of  $B(X)$  generated by any two commuting hermitian operators and the identity operator always has property SN. From a result of C. A. McCarthy's [14], G. Lumer [13] has deduced the existence of a hermitian operator  $L \in B(X)$ , where  $X$  is reflexive but non-Hilbert, whose powers are not all hermitian. In this space  $X$ , the closed subalgebra of  $B(X)$  generated by two commuting hermitian operators and the identity operator need not have property SN.

2.4.2 Using the operator  $L$  in 2.4.1 let us form the family  $L + \lambda I$ , where  $\lambda$  is real and  $I$  is the identity operator. Each member of this family is pseudonormal but not normal.

2.4.3 It is clear from 2.3 that a pseudonormal operator which is such that the closed subalgebra of  $B(X)$  generated by its components and the identity operator has property



$SN$  , is normal. Thus on a Hilbert space pseudonormal and normal operators coincide.

## 2.5 Some properties of normal and pseudonormal operators

2.5.1 Let  $X$  be reflexive. If  $T \in B(X)$  is normal, then  $T$  is spectral of scalar type and so has the spectral representation

$$T = \int_{\sigma(T)} \lambda P(d\lambda) ,$$

where  $P(.)$  is the resolution of the identity for  $T$  .  
If  $T$  is spectral of scalar type, then there exists a norm on  $X$  equivalent to the given norm relative to which  $T$  is normal.

### Proof:

Since  $T$  is normal, we have  $T = u + iv$  where  $uv = vu$  and  $u^m v^n$  are hermitian operators for all  $m, n = 0, 1, 2, \dots$  . Hence by 1.11.3,  $T$  is spectral of type and so

$$T = \int_{\sigma(T)} \lambda P(d\lambda) .$$

Now suppose  $T$  is spectral of scalar type. Then by 1.11.2 there exist operators  $u, v \in B(X)$  such that  $T = u + iv$  ,  $uv = vu$  and relative



to some norm on  $X$  equivalent to the given norm,  $u^m v^n$  are hermitian operators for  $m, n = 0, 1, 2, \dots$ . Relative to this norm,  $T$  is normal.

2.5.2 If  $T \in B(X)$  is normal, then so are  $\lambda T$  and  $T + \lambda$  for all complex  $\lambda$ .

Proof:

Let  $T = u + iv$ , where  $uv = vu$  and  $u^m v^n$  are hermitian operators for  $m, n = 0, 1, 2, \dots$ .

Let  $\lambda = \lambda_1 + i\lambda_2$ , where  $\lambda_1$  and  $\lambda_2$  are real. Then

$$\begin{aligned}\lambda T &= (\lambda_1 + i\lambda_2)(u + iv) \\ &= (\lambda_1 u - \lambda_2 v) + i(\lambda_2 u + \lambda_1 v) \quad .\end{aligned}$$

Since  $\lambda_1, \lambda_2$  are real,  $(\lambda_1 u - \lambda_2 v)^M (\lambda_2 u + \lambda_1 v)^N$  is a polynomial in  $u$  and  $v$  with real coefficients for  $M, N = 0, 1, 2, \dots$ . Since every term  $u^m v^n$  is hermitian it follows that  $(\lambda_1 u - \lambda_2 v)^M (\lambda_2 u + \lambda_1 v)^N$  is also hermitian.

$$\begin{aligned}\text{Now } &(\lambda_1 u - \lambda_2 v)(\lambda_2 u + \lambda_1 v) \\ &= \lambda_1 \lambda_2 u^2 + \lambda_1^2 vu - \lambda_2^2 uv - \lambda_2 \lambda_1 v^2, \text{ since } uv = vu \\ &= (\lambda_2 u + \lambda_1 v)(\lambda_1 u - \lambda_2 v) \quad .\end{aligned}$$

Hence  $\lambda T$  is normal.





$$\begin{aligned} T + \lambda &= (u + iv) + (\lambda_1 + i\lambda_2) \\ &= (u + \lambda_1) + i(v + \lambda_2) \quad . \end{aligned}$$

For  $M, N = 0, 1, 2, \dots$ ,  $(u + \lambda_1)^M (v + \lambda_2)^N$  is a polynomial in  $u$  and  $v$  with real coefficients and since every term  $u^m v^n$  is hermitian for  $m, n = 0, 1, 2, \dots$ , it follows that  $(u + \lambda_1)^M (v + \lambda_2)^N$  is also hermitian for all  $M, N = 0, 1, 2, \dots$ .

$$\begin{aligned} \text{Now} \quad & (u + \lambda_1)(v + \lambda_2) \\ &= vu + \lambda_2 u + \lambda_1 v + \lambda_1 \lambda_2 \quad , \quad \text{since } uv = vu \\ &= (v + \lambda_2)(u + \lambda_1) \quad . \end{aligned}$$

Hence  $T + \lambda$  is normal. The result 2.5.2 is easily seen to hold for pseudonormal operators  $T \in B(X)$ .

2.5.3 If  $T \in B(X)$  is normal, then  $T^n$  is normal and spectraloid for all non-negative integral  $n$ .

Proof:

Let  $T = u + iv$  where  $uv = vu$  and  $u^s v^t$  are hermitian operators for all  $s, t = 0, 1, 2, \dots$ .

$$\begin{aligned} T^n &= (u + iv)^n \\ &= P + iQ \quad , \end{aligned}$$



where  $P$  and  $Q$  are commuting polynomials in  $u$  and  $v$  with real coefficients. Also  $P^M Q^N$  is a polynomial in  $u$  and  $v$  with real coefficients for  $M, N = 0, 1, 2, \dots$ . Since each term  $u^s v^t$  is hermitian for  $s, t = 0, 1, 2, \dots$ , it follows that  $P^M Q^N$  is hermitian for  $M, N = 0, 1, 2, \dots$ . Thus  $T^n$  is normal. Now  $T^n = P + iQ$  belongs to the closed subalgebra of  $B(X)$  generated by  $u, v$  and the identity operator  $I$ , and since by 2.3 this closed subalgebra has property SN, it follows that  $T^n$  is spectro-normaloid.

2.5.4 If  $T$  is normal and  $X$  is reflexive then there exist operators  $R, S \in B(X)$  such that

$$(i) \quad T = RS = SR$$

(ii)  $\sigma(R)$  is a set of non-negative real numbers  
and  $\sigma(S)$  is a subset of the unit circle.

Proof:

By 2.5.1, since  $T$  is normal and  $X$  is reflexive,  $T$  is spectral of scalar type. Hence by 1.12 the result follows.

2.5.5 Lemma If  $X$  is reflexive, then an operator  $T \in B(X)$  is hermitian if and only if  $T'$ , the adjoint of  $T$ , is hermitian.



Proof:

Let  $T \in B(X)$  be hermitian. Then  $W(T) = \{[Tx, x] : \|x\| = 1 \text{ and } x \in X\}$  is real. Since  $X'$ , the conjugate of  $X$ , is a Banach space we may introduce a semi-inner product on it. By the Hahn-Banach theorem in  $X'$ , there exists for an arbitrary but fixed  $f \in X'$ , an  $F_f \in X''$  such that

$$2.5.5.1 \quad F_f(f) = \|f\|^2 \quad \text{and} \quad \|F_f\| = \|f\| \quad .$$

Thus we may define a semi-inner product  $[\cdot, \cdot]'$  on  $X'$  by requiring

$$2.5.5.2 \quad [g, f]' = F_f(g) \quad , \quad g \in X' \quad .$$

Let  $\|f\| = 1$  and consider  $[T'f, f]'$ . By 2.5.5.2 we have

$$[T'f, f]' = F_f(T'f) \quad .$$

Since  $X$  is reflexive, there exists a unique  $x_f \in X$  such that

$$F_f(T'f) = (T'f)(x_f)$$

and

$$\|x_f\| = \|F_f\| \quad .$$

Thus,

$$\begin{aligned} 2.5.5.3 \quad [T'f, f]' &= F_f(T'f) \\ &= (T'f)(x_f) \\ &= f(Tx_f) \quad , \end{aligned}$$



by the definition of  $T'$  .

Now  $x_f$  corresponds to  $F_f$  in the canonical mapping. Hence  $F_f(f) = f(x_f)$  . Thus by 2.5.5.1,

$$f(x_f) = F_f(f) = \|f\|^2 = \|F_f\|^2 = \|x_f\|^2$$

and

$$\|f\| = \|F_f\| = \|x_f\| \quad .$$

Hence

$$f(x_f) = \|x_f\|^2$$

and

$$\|f\| = \|x_f\| \quad .$$

Let us now define a semi-inner product  $[\cdot, \cdot]_1$  on  $X$  where we use the function  $f$  at the point  $x_f$  , that is such that

$$[y, x_f]_1 = f(y) \quad \text{for all } y \in X .$$

Then by 2.5.5.3, we have

$$\begin{aligned} [T'f, f]' &= f(Tx_f) \\ &= [Tx_f, x_f]_1 \end{aligned}$$

Since  $\|x_f\| = \|f\| = 1$  , it follows that  $[Tx_f, x_f]_1 \in W_1(T)$  , the numerical range of  $T$  determined by the semi-inner product  $[\cdot, \cdot]_1$  on  $X$  . Since  $W(T)$  is real, we have by 1.3.2 that any other determination  $W_1(T)$  of the numerical range of  $T$  is real.





In particular,  $[Tx_f, x_f]_1$  is real.

Hence  $[T'f, f]'$  is real. Since  $f \in X'$  is arbitrary with  $\|f\| = 1$ , it follows that the numerical range of  $T'$  determined by the semi-inner product  $[\cdot, \cdot]'$  on  $X'$  is real. Thus  $T'$  is a hermitian operator on  $X'$ .

Suppose now that  $T'$  is a hermitian operator on  $X'$ . Then by what has been proved  $T''$  is a hermitian operator on  $X'' = X$ ,  $X$  being reflexive. Also  $T'' = T$ . Thus  $T$  is a hermitian operator on  $X$ .

Remark.

It follows from the property  $\|T\| = \|T'\|$  and 1.4.1 that lemma 2.5.5 holds for non-reflexive Banach spaces also.

2.5.6 If  $X$  is reflexive, then an operator  $T \in B(X)$  is normal if and only if  $T'$ , the adjoint of  $T$ , is normal.

Proof:

Suppose  $T$  is normal and let  $T = u + iv$  where  $uv = vu$  and  $u^m v^n$  are hermitian for all  $m, n = 0, 1, 2, \dots$ . Now  $T' = u' + iv'$  and

$$(uv)' = v'u' = (vu)' = u'v'.$$

Since  $u^m v^n$  are all hermitian we have by 2.5.5 that

$$(u^m v^n)' = u'^m v'^n$$



are hermitian for all  $m, n = 0, 1, 2, \dots$ . Hence  $T'$  is normal.

Now suppose that  $T'$  is normal. Then by the immediately preceding argument we have  $T''$  normal. This implies, since  $T'' = T$ , that  $T$  is normal.

The result 2.5.6 clearly holds for pseudonormal operators  $T \in B(X)$ .

### 2.5.7 Fuglede's theorem

Let  $X$  be reflexive. Let  $T \in B(X)$  be normal. Then any operator  $B \in B(X)$  which commutes with  $T$ , commutes with  $P(\Delta)$  for all  $\Delta \in \mathcal{S}$  (see 1.9), where  $P(\cdot)$  is the resolution of the identity for  $T$ , (see [8]).

#### Proof:

Since  $T$  is normal, it follows from 2.5.1 that it is spectral (of scalar type). Hence the result follows from 1.9.9.

2.5.8 Let  $X$  be reflexive. Let  $S, T \in B(X)$  be normal and suppose that  $S = u_s + iv_s$ ,  $T = u_T + iv_T$  where  $u_s v_s = v_s u_s$ ,  $u_T v_T = v_T u_T$  and  $u_s^m v_s^n$ ,  $u_T^m v_T^n$  are hermitian operators for all  $m, n = 0, 1, 2, \dots$ . If  $ST = TS$ , then  $S + T$  is pseudonormal. If in addition  $(u_s^m v_s^n) (u_T^{m'} v_T^{n'})$  are hermitian for  $m, n, n', m' = 0, 1, 2, \dots$ , then  $S + T$  and  $ST$  are both normal.



Proof:

Since  $S$  and  $T$  are normal, they are spectra<sup>1</sup> (of scalar type) by 2.5.1 and so have resolutions of the identity  $P_S(\cdot)$  and  $P_T(\cdot)$  respectively. By Fuglede's theorem (2.5.7), since  $S$  is normal and commutes with  $T$ ,  $T$  commutes with  $P_S(\cdot)$ . Since  $P_S(\cdot)$  commutes with  $T$  and  $T$  is normal, it follows similarly that  $P_S(\cdot)$  commutes with  $P_T(\cdot)$ . Thus

$$2.5.8.1 \quad P_S(\cdot) P_T(\cdot) = P_T(\cdot) P_S(\cdot) \quad .$$

Now by the uniqueness (2.1) of  $u_s, v_s, u_T, v_T$  and the spectral representations (2.5.1):

$$T = \int_{\sigma(T)} \lambda P_T(d\lambda) , \quad s = \int_{\sigma(s)} \mu P_S(d\mu)$$

we have

$$u_s = \int_{\sigma(s)} \operatorname{Re} \lambda P_S(d\lambda) , \quad v_s = \int_{\sigma(s)} \operatorname{Im} \lambda P_S(d\lambda)$$

$$u_T = \int_{\sigma(T)} \operatorname{Re} \mu P_T(d\mu) , \quad v_T = \int_{\sigma(T)} \operatorname{Im} \mu P_T(d\mu)$$

where " $\operatorname{Re} \lambda$ " and " $\operatorname{Im} \lambda$ " denote the real and imaginary parts of  $\lambda$  respectively.



Hence

$$u_s u_T = \int_{\sigma(s)} \operatorname{Re} \lambda P_s(d\lambda) \times \int_{\sigma(T)} \operatorname{Re} \mu P_T(d\mu) \quad .$$

Thus if  $\pi_1 = (\Delta_1, \Delta_2, \dots, \Delta_n)$  and  $\pi_2 = (\delta_1, \delta_2, \dots, \delta_m)$  are partitions of  $\sigma(s)$  and  $\sigma(T)$  respectively with norms,  $\|\pi_1\|$ ,  $\|\pi_2\|$  (see 1.9.10), then

$$\begin{aligned} u_s u_T &= \lim_{\|\pi_1\| \rightarrow 0} \sum_{i=1}^n \operatorname{Re} \lambda_i P_s(\Delta_i) \times \lim_{\|\pi_2\| \rightarrow 0} \sum_{j=1}^m \operatorname{Re} \mu_j P_T(\delta_j) \\ &= \lim_{\|\pi_1\| \rightarrow 0} \lim_{\|\pi_2\| \rightarrow 0} \sum_{i,j=1}^{n,m} (\operatorname{Re} \lambda_i)(\operatorname{Re} \mu_j) P_s(\Delta_i) P_T(\delta_j) \quad . \end{aligned}$$

Hence by 2.5.8.1, we have

$$\begin{aligned} u_s u_T &= \lim_{\|\pi_1\| \rightarrow 0} \lim_{\|\pi_2\| \rightarrow 0} \sum_{i,j=1}^{n,m} (\operatorname{Re} \lambda_i)(\operatorname{Re} \mu_j) P_T(\delta_j) P_s(\Delta_i) \\ &= \lim_{\|\pi_2\| \rightarrow 0} \sum_{j=1}^m \operatorname{Re} \mu_j P_T(\delta_j) \times \lim_{\|\pi_1\| \rightarrow 0} \sum_{i=1}^n \operatorname{Re} \lambda_i P_s(\Delta_i) \\ &= \int_{\sigma(T)} \operatorname{Re} \mu P_T(d\mu) \times \int_{\sigma(s)} \operatorname{Re} \lambda P_s(d\lambda) \\ &= u_T u_s \quad . \end{aligned}$$





Similarly,  $u_s v_T = v_T u_s$ ,  $v_s u_T = u_T v_s$ ,  $v_s v_T = v_T v_s$ .

Thus we have

$$2.5.8.2 \quad u_s u_T = u_T u_s$$

$$u_s v_T = v_T u_s$$

$$v_s u_T = u_T v_s$$

$$v_s v_T = v_T v_s$$

Now  $S + T = (u_s + u_T) + i(v_s + v_T)$ . Thus by 2.5.8.2 we have

$$\begin{aligned} & (u_s + u_T)(v_s + v_T) \\ &= u_s v_s + u_s v_T + u_T v_s + u_T v_T \\ &= v_s u_s + v_T u_s + v_s u_T + v_T u_T \\ &= (v_s + v_T)(u_s + u_T) \end{aligned}$$

Since  $u_s, u_T, v_s, v_T$  are hermitian operators, so are  $u_s + u_T$  and  $v_s + v_T$ . Thus  $S + T$  is pseudonormal.

For all  $M, N = 0, 1, 2, \dots$ ,  $(u_s + u_T)^M (v_s + v_T)^N$  is the sum of real multiples of terms of the form

$$\begin{aligned} & u_s^m u_T^n v_s^{m'} v_T^{n'} \\ &= (u_s^m v_s^{m'}) (u_T^n v_T^{n'}). \end{aligned}$$



Since by hypothesis  $(u_s^m v_s^{m'})(u_T^n v_T^{n'})$  is hermitian for all  $m, m', n, n' = 0, 1, 2, \dots$ , it follows that  $(u_s + u_T)^M (v_s + v_T)^N$  is also hermitian for all  $M, N = 0, 1, 2, \dots$ . Hence  $S + T$  is normal.

$$ST = (u_s u_T - v_s v_T) + i(u_s v_T + v_s u_T)$$

Also  $(u_s u_T - v_s v_T)$  and  $(u_s v_T + v_s u_T)$  commute on account of 2.5.8.2 .

Now for  $M, N = 0, 1, 2, \dots$ ,  $(u_s u_T - v_s v_T)^M (u_s v_T + v_s u_T)^N$  is the sum of real multiples of terms of the form

$$(u_s^m v_s^n)(u_T^{m'} v_T^{n'}) .$$

Thus by our hypothesis,

$$(u_s u_T - v_s v_T)^M (u_s v_T + v_s u_T)^N$$

is hermitian for all  $M, N = 0, 1, 2, \dots$ . Hence  $ST$  is normal.

2.5.9      Lemma      Let  $T \in B(X)$  . If  $T = u + iv$  where  $u$  and  $v$  are hermitian operators in  $B(X)$  and  $M$  is one of a pair  $(M, N)$  of subspaces of  $X$  which reduces  $T$  completely, then

$$T|_M = u|_M + iv|_M ,$$

where  $u|_M$  and  $v|_M$  are both hermitian operators in  $B(M)$  .



Proof:

By 1.5,

$$X = M \oplus N .$$

Also  $M$  and  $N$  are both invariant under  $T$  and so  $T|_M$  and  $T|_N$  are both defined. For arbitrary  $x \in X$ , we can write

$$x = x_1 + x_2 , \quad \text{where } x_1 \in M , \quad x_2 \in N .$$

Hence  $Tx = Tx_1 + Tx_2$  and so, since  $x_1 \in M$  and  $x_2 \in N$  we have

$$ux + ivx = T|_M x_1 + T|_N x_2 .$$

If  $x \in M$ , then  $x_1 = x$  and  $x_2 = 0$  and so

$$\begin{aligned} ux + ivx &= T|_M x + T|_N 0 \\ &= T|_M x . \end{aligned}$$

Thus

$$u|_M x + iv|_M x = T|_M x .$$

This holds for arbitrary  $x \in M$ , and so

$$T|_M = u|_M + iv|_M .$$

Now  $W\{u|_M\}$  and  $W\{v|_M\}$  are subsets respectively of  $W(u)$  and  $W(v)$  and since both of these are real on account of the hermiticity of  $u$



and  $v$ , it follows that  $W\{u|_M\}$  and  $W\{v|_M\}$  are both real.

Thus  $T|_M = u|_M + iv|_M$ , where  $u|_M$  and  $v|_M$  are hermitian operators in  $B(M)$ .

2.5.10 If  $T \in B(X)$  is normal and  $M$  is one of a pair of subspaces of  $X$  which reduces  $T$  completely, then  $T|_M$  is also normal.

Proof:

Let  $T = u + iv$ , where  $uv = vu$  and  $u^m v^n$  are hermitian operators for  $m, n = 0, 1, 2, \dots$ .

By 2.5.9, we have

$$T|_M = u|_M + iv|_M, \text{ where } u|_M \text{ and } v|_M$$

are hermitian operators in  $B(M)$ . Since  $uv = vu$ , we have

$$(uv)|_M = (vu)|_M.$$

$$\text{Thus } u|_M v|_M = v|_M u|_M.$$

Also  $(u^m v^n)|_M = (u|_M)^m (v|_M)^n$  are hermitian for  $m, n = 0, 1, 2, \dots$ .

Hence  $T|_M$  is normal.

2.5.11 The limit in the uniform operator topology of a sequence of normal operators is normal.





Proof:

Let  $T_n$  be normal for each  $n$  and suppose that  $T_n \rightarrow T$ .

We can write  $T_n = u_n + iv_n$  where  $u_n v_n = v_n u_n$  and  $u_n^s v_n^t$  are hermitian for all  $s, t = 0, 1, 2, \dots$ .

By an argument similar to that used in the proof of lemma 2.3.3,

$u_n$  and  $v_n$  converge to  $u_o$  and  $v_o$  which commute. Also

$T = u_o + iv_o$ . Since  $u_n^s v_n^t$  is hermitian, it follows that

$$u_o^s v_o^t = \lim_{n \rightarrow \infty} u_n^s v_n^t$$

is hermitian for all  $s, t = 0, 1, 2, \dots$ . Thus  $T$  is normal.



## CHAPTER III

### HYPONORMAL OPERATORS

#### 3.1 Definition

An operator  $T \in B(X)$  will be called hyponormal if and only if .

3.1.1  $T$  can be written in the form  $T = u + iv$  , where  $u$  and  $v \in B(X)$  and are hermitian operators.

3.1.2  $W\{i(uv - vu)\} \geq 0$  , where  $W$  is the numerical range with respect to some semi-inner product on  $X$  .

3.1.3 For any subspace  $M$  of  $X$  which is one of a pair that reduces  $T$  completely, we have  $(T - \lambda I)|_M$  is spectronormaloid for  $\lambda = 0$  or  $\lambda \in \sigma(T)$  the spectrum of  $T$  .

#### 3.2 Some comments on the definition

3.2.1 The condition 3.1.2 always makes sense because I. Vidav [19, p. 123] has shown that if  $u$  and  $v$  are hermitian then so is  $i(uv - vu)$ .

3.2.2 Since a Banach space can in general be made into a semi-



inner product space by the introduction of any one of infinitely many semi-inner products on  $X$ , we must show that the condition 3.1.2 is independent of the semi-inner product we use on  $X$ . Let  $T = u + iv$ , where  $u$  and  $v$  are hermitian operators in  $B(X)$ , satisfy 3.1.2 with respect to some semi-inner product  $[\cdot, \cdot]$  on  $X$ . Then

$$W\{i(uv-vu)\} = \{[i(uv-vu)x, x] : \|x\| = 1\} \geq 0.$$

Let  $[\cdot, \cdot]_1$  be any other semi-inner product on  $X$  and let

$$W_1\{i(uv - vu)\} = \{[i(uv - vu)x, x]_1 : \|x\| = 1\}.$$

By 1.3.1, we have

$$\mathcal{C}W_1\{i(uv - vu)\} = \mathcal{C}W\{i(uv - vu)\} \geq 0,$$

where  $\mathcal{C}$  denotes convex hull.

Hence  $W_1\{i(uv - vu)\} \geq 0$ .

Thus if  $T$  satisfies 3.1.2 for one semi-inner product on  $X$ , then it satisfies it for any other semi-inner product.

3.2.3 If  $X$  is a Hilbert space, the condition 3.1.2 is equivalent to the condition  $T^*T - TT^* \geq 0$ .



Proof:

Since  $X$  is a Hilbert space, there is a unique semi-inner product on  $X$  (see 1.2.5) namely the inner product. Thus condition 3.1.2 is now

$$3.2.3.1 \quad \{(i[uv - vu]x, x) : \|x\| = 1\} \geq 0 .$$

Let  $z \in X$  be arbitrary. If  $z = 0$ , then we have

$$3.2.3.2 \quad (i[uv - vu]z, z) = 0 .$$

If  $z \neq 0$ , it is easily verified that 3.2.3.1 leads to

$$(i[uv - vu]z, z) \geq 0 .$$

Thus by 3.2.3.2 we have

$$(i[uv - vu]x, x) \geq 0 \quad \text{for all } x \in X .$$

Conversely if  $(i[uv - vu]x, x) \geq 0$  for all  $x \in X$ , then

$$(i[uv - vu]x, x) \geq 0 \quad \text{for all } x \text{ such that } \|x\| = 1 .$$

That is  $\{(i[uv - vu]x, x) : \|x\| = 1\} \geq 0$ . Thus if  $X$  is a Hilbert space, the condition 3.1.2 is equivalent to

$$3.2.3.3 \quad \{(i[uv - vu]x, x) : x \in X\} \geq 0 .$$

Since  $X$  is a Hilbert space, every  $T \in B(X)$  can be written  $T = u + iv$ ,





where  $u$  and  $v \in B(X)$  and are hermitian operators. Then  $T^* = u - iv$ . Hence  $T^*T - TT^* = 2i(uv - vu)$ . Thus since  $T$  satisfies 3.1.2 if and only if it satisfies 3.2.3.3, we have

$$\{([T^*T - TT^*]x, x) : x \in X\} \geq 0 .$$

Hence  $T^*T - TT^* \geq 0$ .

3.2.4 If  $X$  is a Hilbert space, then condition 3.1.2 implies condition 3.1.3.

Proof:

By 3.2.3, the condition 3.1.2 is equivalent to  $T^*T - TT^* \geq 0$ . This condition in a Hilbert space is equivalent to the hyponormality of  $T$  according to the usual Hilbert space definition of hyponormality, (see 1.14.1).

By 1.14.2,  $T - \lambda$  is thus hyponormal for all complex  $\lambda$  and hence in particular for  $\lambda = 0$  or  $\lambda \in \sigma(T)$ .

If  $M$  is one of a pair of subspaces of  $X$  which reduces  $T$  completely then  $M$  is invariant under  $T - \lambda$  and since  $T - \lambda$  is hyponormal, we have by 1.14.1 that  $(T - \lambda)|_M$  is hyponormal. Hence by 1.14.3,

$$\|(T - \lambda)|_M\| = r[(T - \lambda)|_M] .$$

The condition 3.1.2 thus implies 3.1.3.



3.2.5 If  $X$  is a Hilbert space, then the definition of hyponormality in 3.1 is equivalent to the usual Hilbert space definition.

Proof:

This follows readily from 3.2.3 and 3.2.4.

### 3.3 Some properties of hyponormal operators

3.3.1 If  $T \in B(X)$  and is hyponormal, then so is  $\lambda T$  for all complex  $\lambda$ . If  $\lambda \in \sigma(T)$ , then  $T - \lambda$  is also hyponormal.

Proof:

Let  $T = u + iv$ , where  $u$  and  $v$  are hermitian operators in  $B(X)$  and  $W\{i(uv - vu)\} \geq 0$ .

Let  $\lambda = \lambda_1 + i\lambda_2$ , where  $\lambda_1$  and  $\lambda_2$  are real. Then

$$\begin{aligned}\lambda T &= (\lambda_1 + i\lambda_2)(u + iv) \\ &= (\lambda_1 u - \lambda_2 v) + i(\lambda_1 v + \lambda_2 u) .\end{aligned}$$

Since  $u$  and  $v$  are hermitian operators, so are  $\lambda_1 u - \lambda_2 v$  and  $\lambda_1 v + \lambda_2 u$ . Now



$$\begin{aligned}
 & W\{i[(\lambda_1 u - \lambda_2 v)(\lambda_1 v + \lambda_2 u) - (\lambda_1 v + \lambda_2 u)(\lambda_1 u - \lambda_2 v)]\} \\
 &= W\{i[\lambda_1^2 + \lambda_2^2](uv - vu)\} \\
 &= (\lambda_1^2 + \lambda_2^2) W\{i(uv - vu)\} \geq 0 .
 \end{aligned}$$

Since  $T$  is hyponormal, we have by definition that if  $M$  is any one of a pair of subspaces that reduces  $T$  completely, then

$$\|(T - \lambda_o)|_M\| = r\{(T - \lambda_o)|_M\} \quad \text{for } \lambda_o = 0 \text{ or } \lambda_o \in \sigma(T) .$$

Let  $\mu = 0$  or  $\mu \in \sigma(\lambda T)$  .

Then we can write  $\mu = \lambda \lambda_o$  . Hence

$$\begin{aligned}
 r\{(\lambda T - \mu)|_M\} &= r\{(\lambda T - \lambda \lambda_o)|_M\} \\
 &= |\lambda| \, r\{(T - \lambda_o)|_M\} \\
 &= |\lambda| \cdot \|(T - \lambda_o)|_M\| \\
 &= \|(\lambda T - \lambda \lambda_o)|_M\| \\
 &= \|(\lambda T - \mu)|_M\| .
 \end{aligned}$$

Thus  $\lambda T$  is hyponormal.

Now  $T - \lambda = (u - \lambda_1) + i(v - \lambda_2)$  . Clearly  $u - \lambda_1$  and  $v - \lambda_2$  are hermitian operators and



$$\begin{aligned} & W\{i[(u-\lambda_1)(v-\lambda_2) - (v-\lambda_2)(u-\lambda_1)]\} \\ & = W\{i(uv - vu)\} \geq 0 \end{aligned}$$

Let  $\mu = 0$  or  $\mu \in \sigma(T - \lambda)$ . Then since  $\lambda \in \sigma(T)$  we can write

$$\mu = \lambda_0 - \lambda, \quad \text{where } \lambda_0 \in \sigma(T).$$

$$\begin{aligned} \text{Thus } [(T - \lambda) - \mu]|_M &= [T - \lambda - (\lambda_0 - \lambda)]|_M \\ &= (T - \lambda_0)|_M. \end{aligned}$$

Hence, since  $T$  is hyponormal and  $\lambda_0 \in \sigma(T)$ , we have by 3.1.3 that

$$r\{[(T - \lambda) - \mu]|_M\} = \|[(T - \lambda) - \mu]|_M\|$$

for  $\mu = 0$  or  $\mu \in \sigma(T - \lambda)$ .

This shows that  $T - \lambda$  is hyponormal.

3.3.2 If  $T \in B(X)$  and is normal, then it is hyponormal.

Proof:

Let  $T = u + iv$  where  $uv = vu$  and  $u^m v^n$  are hermitian operators for  $m, n = 0, 1, 2, \dots$ .

$$\text{Then } W\{i(uv - vu)\} = W\{0\} = 0.$$

Since  $T$  is normal, so is  $T + \lambda$  for all complex  $\lambda$  (see 2.5.2). In particular  $T - \lambda_0$  is normal for all  $\lambda_0 = 0$  or  $\lambda_0 \in \sigma(T)$ . Thus





by 2.5.10, if  $M$  is any one of a pair of subspaces of  $X$  which reduces  $T$  completely, then  $(T - \lambda_0)|_M$  is also normal. Hence by 2.5.3,  $(T - \lambda_0)|_M$  is spectro-normaloid.

Thus  $T$  is hyponormal.

3.3.3 If  $T \in B(X)$  and is hyponormal and  $M$  is any one of a pair of subspaces of  $X$  which reduces  $T$  completely, then  $T|_M$  is hyponormal.

Proof:

Since  $T$  is hyponormal, we can write  $T = u + iv$ , where  $u$  and  $v$  are hermitian operators in  $B(X)$  such that

$$W\{i(uv - vu)\} \geq 0.$$

Since  $M$  is one of a pair of subspaces that reduces  $T$  completely, we have by lemma 2.5.9, that

$$T|_M = u|_M + iv|_M \quad \text{where} \quad u|_M \quad \text{and} \quad v|_M$$

are hermitian operators in  $B(M)$ .

$$\begin{aligned} \text{Now} \quad & W\{i(u|_M v|_M - v|_M u|_M)\} \\ &= W\{i(uv - vu)|_M\} \geq 0 \end{aligned}$$

since  $W\{i(uv - vu)|_M\}$  is a subset of  $W\{i(uv - vu)\}$ .

Let  $N \subset M$  be any one of a pair of subspaces of  $M$  which reduces  $T|_M$



completely. Then clearly  $N$  is one of a pair of subspaces of  $X$  which reduces  $T$  completely. Also for any  $\lambda$ ,

$$(T - \lambda)|_N = \{(T - \lambda)|_M\}|_N .$$

Since  $T$  is hyponormal,  $(T - \lambda)|_N$  is spectro-normaloid for  $\lambda = 0$  or  $\lambda \in \sigma(T)$ . Hence  $(T|_M - \lambda)|_N$  is spectro-normaloid for  $\lambda = 0$  or  $\lambda \in \sigma(T)$ . Thus in particular  $(T|_M - \lambda)|_N$  is spectro-normaloid for  $\lambda = 0$  or  $\lambda \in \sigma(T|_M) \subseteq \sigma(T)$ . Thus  $T|_M$  is hyponormal.

3.3.4 The only generalised quasi-nilpotent hyponormal operator in  $B(X)$  is the zero operator.

Proof:

Let  $T$  be a generalised quasi-nilpotent hyponormal operator in  $B(X)$ . Then  $r(T) = 0$ .

Since  $T$  is hyponormal, we have by 3.1.3 that if  $M$  is any one of a pair of subspaces of  $X$  which reduces  $T$  completely then  $(T - \lambda)|_M$  is spectro-normaloid for  $\lambda = 0$  or  $\lambda \in \sigma(T)$ . Let us choose  $M = X$  and  $\lambda = 0$ .

Then we have

$$\|T\| = r(T) = 0 .$$

Thus  $T = 0$ .



3.3.5 Let  $T$  be a hyponormal operator in  $B(X)$ . If  $\lambda_o$  is an isolated point of  $\sigma(T)$ , then  $\lambda_o \in P\sigma(T)$ . Also the subspace  $M(\lambda_o) = \{x \in X : Tx = \lambda_o x\}$  is one of a pair which reduces  $T$  completely, and  $T|_{M(\lambda_o)}$  is normal.

Proof:

Since  $\lambda_o$  is an isolated part of  $\sigma(T)$ , it follows by the decomposition theorem (see 1.6) that there exists a subspace  $M_{\{\lambda_o\}}$  of  $X$  which is one of a pair which reduces  $T$  completely and is such that

$$\sigma(T|_{M_{\{\lambda_o\}}}) = \{\lambda_o\}.$$

Also by 3.3.3,  $T|_{M_{\{\lambda_o\}}}$  is hyponormal. Hence  $T|_{M_{\{\lambda_o\}}} - \lambda_o$  is hyponormal by 3.3.1 since

$$\lambda_o \in \sigma(T|_{M_{\{\lambda_o\}}}).$$

Now

$$\begin{aligned} & \sigma(T|_{M_{\{\lambda_o\}}} - \lambda_o) \\ &= \{\sigma(T|_{M_{\{\lambda_o\}}}) - \lambda_o\} \\ &= \{\lambda_o - \lambda_o\} \\ &= \{0\}. \end{aligned}$$



Hence

$$r\{T|_{M_{\{\lambda_o\}}} - \lambda_o\} = 0 \quad .$$

Thus  $T|_{M_{\{\lambda_o\}}} - \lambda_o$  is a generalised quasi-nilpotent hyponormal operator.

Hence by 3.3.4,

$$T|_{M_{\{\lambda_o\}}} - \lambda_o \text{ is the zero operator in } B(M_{\{\lambda_o\}}) \quad .$$

Thus

$$T|_{M_{\{\lambda_o\}}} x = \lambda_o x \text{ for all } x \in M_{\{\lambda_o\}} \quad .$$

Using theorem 1.6 we in fact have,

$$M_{\{\lambda_o\}} = \{x \in X : Tx = \lambda_o x\} = M(\lambda_o) \quad .$$

Since by 1.6  $M_{\{\lambda_o\}}$  is non-trivial, there is some  $x \neq 0$  in  $M(\lambda_o)$  .

Hence  $\lambda_o \in P\sigma(T)$  .

Since  $T$  is hyponormal, we can write  $T = u + iv$  where  $u$  and  $v$  are hermitian operators in  $B(X)$  .

Since  $M(\lambda_o)$  is one of a pair of subspaces which reduces  $T$  completely we have by lemma 2.5.9, that

$$T|_{M(\lambda_o)} = u|_{M(\lambda_o)} + iv|_{M(\lambda_o)}$$

where  $u|_{M(\lambda_o)}$  and  $v|_{M(\lambda_o)}$  are hermitian operators in  $B[M(\lambda_o)]$  .





For  $\|x\| = 1$  and  $x \in M(\lambda_o)$  we have

$$[T|_{M(\lambda_o)} x, x] = [u|_{M(\lambda_o)} x, x] + i[v|_{M(\lambda_o)} x, x] \quad .$$

Hence

$$[\lambda_o x, x] = [u|_{M(\lambda_o)} x, x] + i[v|_{M(\lambda_o)} x, x] \quad , \quad \text{since } T|_{M(\lambda_o)} = \lambda_o \quad .$$

Let  $\lambda_o = \lambda_1 + i\lambda_2$  , where  $\lambda_1$  and  $\lambda_2$  are real. Then

$$[(u|_{M(\lambda_o)} - \lambda_1)x, x] = i[(\lambda_2 - v|_{M(\lambda_o)})x, x] \quad .$$

But  $u|_{M(\lambda_o)} - \lambda_1$  , and  $\lambda_2 - v|_{M(\lambda_o)}$  are both hermitian operators in  $B[M(\lambda_o)]$  and so

$$[(u|_{M(\lambda_o)} - \lambda_1)x, x] \quad \text{and} \quad [(\lambda_2 - v|_{M(\lambda_o)})x, x]$$

are both real.

It follows then that

$$[(u|_{M(\lambda_o)} - \lambda_1)x, x] = 0 = [(\lambda_2 - v|_{M(\lambda_o)})x, x] \quad .$$

But by [2, p. 336],

$$\|u|_{M(\lambda_o)} - \lambda_1\| \leq 4 \sup \{ |\lambda| : \lambda \in W[u|_{M(\lambda_o)} - \lambda_1] \}$$

Hence

$$\|u|_{M(\lambda_o)} - \lambda_1\| = 0 \quad .$$

That is



$$u|_{M(\lambda_o)} = \lambda_1 I|_{M(\lambda_o)} \quad .$$

Similarly,

$$v|_{M(\lambda_o)} = \lambda_2 I|_{M(\lambda_o)} \quad .$$

Hence

$$u|_{M(\lambda_o)} v|_{M(\lambda_o)} = \lambda_1 \lambda_2 I|_{M(\lambda_o)} = v|_{M(\lambda_o)} u|_{M(\lambda_o)} \quad .$$

Also

$$\{u|_{M(\lambda_o)}\}^m \{v|_{M(\lambda_o)}\}^n = \lambda_1^m \lambda_2^n I|_{M(\lambda_o)}$$

is a hermitian operator in  $B[M(\lambda_o)]$  for all  $m, n = 0, 1, 2, \dots$  .

Thus  $T|_{M(\lambda_o)}$  is normal.

3.3.6 Let  $T$  be a non-zero hyponormal operator in  $B(X)$  .

If  $\sigma(T)$  contains exactly one cluster point, then

$T$  is normal.

Proof:

Without loss of generality we may assume that the single cluster point in  $\sigma(T)$  is 0 , for if the cluster point  $\mu_o \neq 0$  , we may consider in place of  $T$  , the operator  $T - \mu_o$  which is also hyponormal by 3.3.1 and has 0 as the single cluster point in its spectrum. Since  $T$  is hyponormal, it is spectro-normaloid by 3.1.3.



Hence

$$\|T\| = r(T) = \max\{|\lambda| : \lambda \in \sigma(T)\} \quad .$$

Let

$$|\lambda_1| = \max\{|\lambda| : \lambda \in \sigma(T)\} \quad .$$

Then

$$|\lambda_1| = \|T\| \neq 0 \quad .$$

Since  $\lambda_1 \neq 0$  , it is an isolated point of  $\sigma(T)$  . Hence by 3.3.5,  $\lambda_1 \in P\sigma(T)$  .

Let  $M_1 = \{x \in X : Tx = \lambda_1 x\}$  .

Then since  $\lambda_1 \in P\sigma(T)$  ,  $M_1$  is non-trivial. Furthermore, by 3.3.5,  $M_1$  is one of a pair of subspaces of  $X$  which reduces  $T$  completely and  $T|_{M_1}$  is normal. Thus if  $N_1$  is the other member of the pair of which  $M_1$  is one, then by 1.5 we also have

$$X = M_1 \oplus N_1 \quad .$$

Hence by 1.6,

$$\sigma(T|_{N_1}) = \sigma(T) - \{\lambda_1\} \quad .$$

Let

$$|\lambda_2| = \max\{|\lambda| : \lambda \in \sigma(T|_{N_1})\} \quad .$$



Then

$$|\lambda_2| \leq |\lambda_1| \quad .$$

Also  $N_1$  is one of a pair of subspaces which reduces  $T$ , and so by 3.3.3,  $T|_{N_1}$  is hyponormal. Thus

$$\|T|_{N_1}\| = r(T|_{N_1}) = |\lambda_2| \quad .$$

Define

$$\begin{aligned} M_2 &= \{x \in N_1 : T|_{N_1} x = \lambda_2 x\} \\ &= \{x \in X : Tx = \lambda_2 x\} \quad . \end{aligned}$$

Applying 3.3.5 again,  $M_2$  is one of a pair of subspace  $(M_2, N_2)$  of  $N_1$  which reduces  $T|_{N_1}$  completely and  $(T|_{N_1})|_{M_2} = T|_{M_2}$  is normal. Thus we have  $X = M_1 \oplus M_2 \oplus N_2$ . Also by 3.3.3,  $(T|_{N_1})|_{N_2} = T|_{N_2}$  is hyponormal and by 1.6,

$$\begin{aligned} \sigma(T|_{N_2}) &= \sigma(T|_{N_1}) - \{\lambda_2\} \\ &= \sigma(T) - \{\lambda_1\} \cup \{\lambda_2\} \quad . \end{aligned}$$

Let

$$|\lambda_3| = \max \{|\lambda| : \lambda \in \sigma(T|_{N_2})\} \quad .$$

Then

$$|\lambda_1| \geq |\lambda_2| \geq |\lambda_3| \quad .$$





Define

$$\begin{aligned} M_3 &= \{x \in N_2 : T|_{N_2} x = \lambda_3 x\} \\ &= \{x \in X : Tx = \lambda_3 x\} \quad , \quad \text{by 1.6} . \end{aligned}$$

We now repeat the argument used twice already.

At the  $n^{\text{th}}$  step we shall obtain  $X = M_1 \oplus M_2 \oplus \dots \oplus M_n \oplus N_n$  ,  
where  $T|_{M_i}$  is normal for each  $i$  and  $T|_{N_n}$  is hyponormal with

$$\|T|_{N_n}\| = |\lambda_{n+1}| = \max \{ |\lambda| : \lambda \in \sigma(T|_{N_n}) \} .$$

Also

$$\|T\| = |\lambda_1| \geq |\lambda_2| \geq \dots \geq |\lambda_{n+1}| > 0 .$$

Thus  $\{\lambda_n\}_{n=1}^{\infty}$  is a sequence in  $\sigma(T)$ , a compact set, and so contains a convergent subsequence. This subsequence can only converge to 0 .

The fact that  $|\lambda_1| \geq |\lambda_2| \geq \dots \geq |\lambda_n| \geq \dots$  then implies that  $\{\lambda_n\}_{n=1}^{\infty}$  converges to 0 , since 0 is the only cluster point in  $\sigma(T)$  . Thus  $\lambda_n \rightarrow 0$  as  $n \rightarrow \infty$  .

Since  $T|_{M_i}$  is normal for each  $i$  , so is  $T|_{M_1 \oplus M_2 \oplus \dots \oplus M_n}$  .

Let us define an operator  $T_n : X \rightarrow X$  where

$$T_n x = \begin{cases} T|_{M_1 \oplus M_2 \oplus \dots \oplus M_n} x & \text{if } x \in M_1 \oplus M_2 \oplus \dots \oplus M_n \\ 0 , & \text{if } x \notin M_1 \oplus M_2 \oplus \dots \oplus M_n . \end{cases}$$

Let us also define an operator  $T_n^{\#} : X \rightarrow X$  where



$$T_n^\# x = \begin{cases} T|_{N_n} x & \text{if } x \in N_n \\ 0, & \text{if } x \notin N_n \end{cases}.$$

Now

$$\begin{aligned} \|T_n^\#\| &= \sup \{ \|T_n^\# x\| : x \in X \text{ and } \|x\| = 1 \} \\ &= \sup \{ \|T|_{N_n} x\| : x \in N_n \text{ and } \|x\| = 1 \} \\ &= \|T|_{N_n}\|. \end{aligned}$$

Also

$$\begin{aligned} \|T_n - T\| &= \sup \{ \|T_n x - Tx\| : x \in X \text{ and } \|x\| = 1 \} \\ &= \sup \{ \|T_n x - Tx\| : x \in M_1 \oplus M_2 \oplus \dots \oplus M_n \text{ and } \|x\| = 1 \} \end{aligned}$$

Let  $x \in X$ . Then  $x = y_1 + y_2$ , where  $y_1 \in M_1 \oplus M_2 \oplus \dots \oplus M_n$  and  $y_2 \in N_n$ . Now  $T_n^\# x = T_n^\# y_1 + T_n^\# y_2$   

$$= T|_{N_n} y_2.$$

Also 
$$\begin{aligned} T_n x &= T_n y_1 + T_n y_2 \\ &= T_n y_1. \end{aligned}$$

Hence

$$\begin{aligned} T_n^\# x + T_n x &= T|_{N_n} y_2 + T_n y_1 \\ &= Ty_2 + Ty_1 \\ &= Tx. \end{aligned}$$



Hence  $T_n x - Tx = -T_n^\# x$  and so

$$\begin{aligned} \|T_n - T\| &= \sup \{ \|T_n^\# x\| : x \in M_1 \oplus M_2 \oplus \dots \oplus M_n \text{ and } \|x\| = 1 \} \\ &\leq \sup \{ \|T_n^\#\| \cdot \|x\| : x \in M_1 \oplus M_2 \oplus \dots \oplus M_n \text{ and } \|x\| = 1 \} \\ &= \|T_n^\#\| \\ &= \|T|_{N_n}\| \\ &= |\lambda_{n+1}|. \end{aligned}$$

Hence

$$\lim_{n \rightarrow \infty} \|T_n - T\| = \lim_{n \rightarrow \infty} |\lambda_{n+1}| = 0.$$

Thus  $T$  is the limit in the uniform operator topology of the sequence  $\{T_n\}_{n=1}^\infty$  of normal operators in  $B(X)$ . It follows by 2.5.11, that  $T$  is normal.

### Corollary I

If  $T$  is a hyponormal completely continuous operator in  $B(X)$  where  $X$  is infinite dimensional and  $P\sigma(T)$  is infinite, then  $T$  is normal.

### Proof:

Since  $T$  is completely continuous and  $X$  is infinite dimensional, we have by theorem 5.5-F and 5.5-G in [18, p. 281-282] that



$$(i) \quad 0 \in \sigma(T)$$

(ii)  $P\sigma(T)$  contains at most a countable set of points with 0 as the only possible cluster point.

Since by hypothesis  $P\sigma(T)$  is infinite, we have by (i) and (ii) and the compactness of  $\sigma(T)$  that 0 is the one cluster point of  $\sigma(T)$ . Thus by 3.3.6,  $T$  is normal.

### Corollary II

If  $T \in B(X)$  is hyponormal, where  $X$  is finite dimensional and  $0 \in \sigma(T)$  then  $T$  is normal.

### Proof:

With the same notation as in the proof of 3.3.6, we have

$$\|T_n - T\| \leq |\lambda_{n+1}| \quad \text{for some } n.$$

Since  $X$  is finite dimensional, every operator  $T \in B(X)$  is completely continuous [see p. 285, problem 1, 18].

Also  $P\sigma(T) = \sigma(T)$ . Since  $0 \in \sigma(T) = P\sigma(T)$ , there is an  $n$  such that  $\lambda_{n+1} = 0$ . For this  $n$ , we have

$$\|T_n - T\| = 0.$$

Hence  $T = T_n$ .

Since  $T_n$  is normal, it follows that  $T$  is also normal.





3.3.7 Not every hyponormal operator is normal.

Proof:

We shall exhibit a counter example. Take  $x = \ell'(n)$ ,  $n \geq 3$ , then the matrix

$$C = \begin{pmatrix} c_{11} & c_{12} & \cdots & c_{1n} \\ c_{21} & & & \\ \cdot & & & \\ \cdot & & & \\ \cdot & & & \\ c_{n1} & & & c_{nn} \end{pmatrix}$$

is a linear operator on  $\ell'(n)$  and

$$\|C\| = \max_{1 \leq \beta \leq n} \left\{ \sum_{\alpha=1}^n |c_{\alpha\beta}| \right\}.$$

Let

$$s_{\beta} = \sum_{\alpha=1}^n c_{\alpha\beta},$$

where the  $c_{\alpha\beta}$  are non-negative real numbers. Consider the matrix

$$M = (c_{\alpha\beta} + i\delta_{\alpha\beta}[s_{\beta} - c_{\beta\beta}]) ,$$



whose  $(\alpha, \beta)^{\text{th}}$  element is  $c_{\alpha\beta} + i\delta_{\alpha\beta}[s_{\beta} - c_{\beta\beta}]$  where  $i = \sqrt{-1}$  and  $\delta_{\alpha\beta}$  is the Kronecker delta.

Let  $t$  be a real, positive, small number. Then if  $I$  is the identity operator on  $\ell'(n)$ , we have

$$\begin{aligned} \|I + it M\| &= \|(\delta_{\alpha\beta} + it[c_{\alpha\beta} + i\delta_{\alpha\beta}\{s_{\beta} - c_{\beta\beta}\}])\| \\ &= \max_{1 \leq \beta \leq n} \left\{ \sum_{\alpha=1}^n |\delta_{\alpha\beta} + it[c_{\alpha\beta} + i\delta_{\alpha\beta}\{s_{\beta} - c_{\beta\beta}\}]| \right\}. \end{aligned}$$

Now

$$\begin{aligned} &\sum_{\alpha=1}^n |\delta_{\alpha\beta} + it[c_{\alpha\beta} + i\delta_{\alpha\beta}(s_{\beta} - c_{\beta\beta})]| \\ &= \sum_{\substack{\alpha=1 \\ \alpha \neq \beta}}^n |itc_{\alpha\beta}| + |1 + it[c_{\beta\beta} + i(s_{\beta} - c_{\beta\beta})]| \\ &= t(s_{\beta} - c_{\beta\beta}) + \sqrt{[1 - t(s_{\beta} - c_{\beta\beta})]^2 + t^2 c_{\beta\beta}^2} \\ &= t(s_{\beta} - c_{\beta\beta}) + \sqrt{1 - 2t(s_{\beta} - c_{\beta\beta}) + t^2\{c_{\beta\beta}^2 + (s_{\beta} - c_{\beta\beta})^2\}} \\ &= t(s_{\beta} - c_{\beta\beta}) + \sqrt{1 - 2t\{s_{\beta} - c_{\beta\beta} - t \frac{L}{2}\}}, \end{aligned}$$

where  $L = c_{\beta\beta}^2 + (s_{\beta} - c_{\beta\beta})^2$ .



$$\begin{aligned}
 &= t(s_{\beta} - c_{\beta\beta}) + 1 + \frac{1}{2} [-2t\{s_{\beta} - c_{\beta\beta} - t \frac{1}{2}\}] \\
 &\quad + \frac{\frac{1}{2} \cdot (\frac{1}{2} - 1)}{2!} 4t^2\{s_{\beta} - c_{\beta\beta} - t \frac{1}{2}\}^2 + \dots
 \end{aligned}$$

$$= 1 + o(t) \quad \text{as } t \rightarrow 0, \quad \text{for all } \beta.$$

Thus

$$\begin{aligned}
 &\max_{1 \leq \beta \leq n} \left\{ \sum_{\alpha=1}^n |\delta_{\alpha\beta} + it[c_{\alpha\beta} + i\delta_{\alpha\beta}(s_{\beta} - c_{\beta\beta})]| \right\} \\
 &= 1 + o(t) \quad \text{as } t \rightarrow 0.
 \end{aligned}$$

Thus  $\|I + it M\| = 1 + o(t)$  as  $t \rightarrow 0$ . Hence by 1.4.1,  $M$  is a hermitian operator in  $\ell^2(n)$ .

Written out in the usual way  $M$  is

$$\begin{array}{ccccc}
 c_{11} + i \sum_{\alpha=2}^n c_{\alpha 1}, & c_{12} & & & \\
 c_{21} & , & c_{22} + i \sum_{\substack{\alpha=1 \\ \alpha \neq 2}}^n c_{\alpha 2}, & & \\
 c_{31} & , & c_{32} & , & \\
 \vdots & & & & \\
 c_{n1} & & & & 
 \end{array}$$



It is easily checked that the matrix obtained from  $M$  and shown below is also a hermitian operator on  $\ell'(n)$  :

$$N = \begin{pmatrix} c_{11} + i \sum_{\alpha=2}^n c_{\alpha 1} & ic_{12} & & \\ ic_{21} & c_{22} + i \sum_{\substack{\alpha=1 \\ \alpha \neq 2}}^n c_{\alpha 2} & & \\ ic_{31} & ic_{32} & & \\ ic_{41} & ic_{42} & & \\ \cdot & & & \\ \cdot & & & \\ \cdot & & & \end{pmatrix}$$

Now we consider the operator  $T = M + iN$  on  $\ell'(n)$  . We have

$$M + iN = \begin{pmatrix} (c_{11} + i \sum_{\alpha=2}^n c_{\alpha 1})(1+i) & & & \\ & (c_{22} + i \sum_{\substack{\alpha=1 \\ \alpha \neq 2}}^n c_{\alpha 2})(1+i) & & \\ & & & \\ & & & \end{pmatrix}$$





where the off-diagonal elements all vanish.

Let us define

$$C_{\beta\beta} = c_{\beta\beta} + i \sum_{\substack{\alpha=1 \\ \alpha \neq \beta}}^n c_{\alpha\beta}$$

$$C_{\alpha\beta} = c_{\alpha\beta} , \quad \text{for } \alpha \neq \beta .$$

Then  $M = (C_{\alpha\beta})$  and

$$N = \begin{pmatrix} C_{11}, & iC_{12}, \\ iC_{21}, & C_{22}, \\ . & \\ . & \\ . & \end{pmatrix}$$

Hence the  $(\alpha, \beta)^{\text{th}}$  element in  $MN$  is

$$\sum_{\substack{r=1 \\ r \neq \beta}}^n iC_{\alpha r} C_{r\beta} + C_{\alpha\beta} C_{\beta\beta} .$$

Also the  $(\alpha, \beta)^{\text{th}}$  element in  $NM$  is



$$\sum_{\substack{r=1 \\ r \neq \alpha}}^n iC_{\alpha r} C_{r\beta} + C_{\alpha\alpha} C_{\alpha\beta} \quad .$$

Hence the  $(\alpha, \beta)^{\text{th}}$  element in  $MN - NM$  is

$$\begin{aligned} i \sum_{r=1}^n C_{\alpha r} C_{r\beta} - iC_{\alpha\beta} C_{\beta\beta} + C_{\alpha\beta} C_{\beta\beta} \\ - i \sum_{r=1}^n C_{\alpha r} C_{r\beta} + iC_{\alpha\alpha} C_{\alpha\beta} - C_{\alpha\alpha} C_{\alpha\beta} \\ = C_{\alpha\beta} (C_{\beta\beta} - C_{\alpha\alpha}) (1-i) \\ = C_{\alpha\beta} (c_{\beta\beta} - c_{\alpha\alpha}) (1-i)^2 \quad . \end{aligned}$$

Thus  $MN \neq NM$  unless  $C_{\alpha\beta} = 0$  or  $c_{\alpha\alpha} = c_{\beta\beta}$  for  $\alpha \neq \beta$  .

Hence as long as we do not satisfy these conditions we can be sure that  $MN \neq NM$  and hence that  $T = M + iN$  is not normal. Now the  $(\alpha, \beta)^{\text{th}}$  element of the Hermitian operator  $i(MN - NM)$  is

$$i(1-i)^2 C_{\alpha\beta} (c_{\beta\beta} - c_{\alpha\alpha}) = 2C_{\alpha\beta} (c_{\beta\beta} - c_{\alpha\alpha}) \quad .$$

Hence if  $x = \{x_1, x_2, \dots, x_n\}$  ,  $y = \{y_1, y_2, \dots, y_n\}$  and  $y = i(MN - NM)x$  , then



$$y_{\alpha} = 2 \sum_{\beta=1}^n c_{\alpha\beta} (c_{\beta\beta} - c_{\alpha\alpha}) x_{\beta} = 2 \sum_{\beta=1}^n c_{\alpha\beta} (c_{\beta\beta} - c_{\alpha\alpha}) x_{\beta} \quad .$$

Now let  $[\cdot, \cdot]$  be a given semi-inner product on  $X$ . Let  $x \in X$  where  $\|x\| = 1$ . Then  $[y, x]$  is a linear function of  $y$ , say  $f(y)$ , such that

$$\|f\| = \|x\| = 1 \quad .$$

Thus we can write [see 18, page 37]

$$\begin{aligned} [y, x] &= f(y) \\ &= \sum_{\alpha=1}^n y_{\alpha} f(\delta^{\alpha}) \quad , \end{aligned}$$

where  $\delta^{\alpha}$  is the unit vector in  $X$  with 1 for its  $\alpha^{\text{th}}$  component.

Hence

$$[i(MN - NM)x, x] = 2 \sum_{\alpha=1}^n f(\delta^{\alpha}) \cdot \sum_{\beta=1}^n c_{\alpha\beta} (c_{\beta\beta} - c_{\alpha\alpha}) x_{\beta} \quad .$$

Let

$$f(\delta^{\alpha}) = F_{\alpha} + i G_{\alpha}$$

$$x_{\beta} = \ell_{\beta} + i m_{\beta} \quad .$$



Then since  $[i(MN-NM)x, x]$  is real, we must have

$$[i(MN-NM)x, x] = 2 \sum_{\alpha=1}^n \sum_{\beta=1}^n c_{\alpha\beta} (c_{\beta\beta} - c_{\alpha\alpha}) (F_{\alpha} \ell_{\beta} - G_{\alpha} m_{\beta}) .$$

Now

$$\begin{aligned} |F_{\alpha} \ell_{\beta} - G_{\alpha} m_{\beta}| &\leq 2 |x_{\beta}| \cdot |f(\delta^{\alpha})| \\ &\leq 2 \|x\| \cdot \|f\| \cdot \|\delta^{\alpha}\| \\ &= 2 . \end{aligned}$$

Hence

$$-2 \leq F_{\alpha} \ell_{\beta} - G_{\alpha} m_{\beta} \leq 2 \quad \text{for all } \alpha \text{ and } \beta .$$

Let

$$m = \inf_{\alpha, \beta} (F_{\alpha} \ell_{\beta} - G_{\alpha} m_{\beta}) .$$

Then  $-2 \leq m \leq 2$  .

(i) Suppose  $m \geq 0$  .

For each  $\beta$  , choose

$$c_{\alpha\beta} = \begin{cases} 0 , & \text{if } \alpha < \beta \\ ar^{\alpha+\beta-2} , & 0 < r < 1, \ a > 0 , \text{ if } \alpha > \beta . \end{cases}$$

With this choice of  $c_{\alpha\beta}$  we have  $MN \neq NM$  and so  $T$  is not normal.

Now we have





$$[i(MN-NM)x, x] = 2 \sum_{\alpha=1}^n \sum_{\beta < \alpha} a^2 r^{\alpha+\beta-2} (r^{2\beta-2} - r^{2\alpha-2}) (F_{\alpha} \ell_{\beta} - G_{\alpha} m_{\beta}) .$$

Since  $0 < r < 1$ , we have

$$r^{2\beta-2} - r^{2\alpha-2} > 0 \quad \text{for } \beta < \alpha .$$

Thus since  $m \leq F_{\alpha} \ell_{\beta} - G_{\alpha} m_{\beta}$  for all  $\alpha$  and  $\beta$ , it follows that

$$\begin{aligned} [i(MN-NM)x, x] &\geq 2 \cdot m \sum_{\alpha=1}^n \sum_{\beta < \alpha} a^2 r^{\alpha+\beta-2} (r^{2\beta-2} - r^{2\alpha-2}) \\ &\geq 0 . \end{aligned}$$

Since  $x$  is arbitrary with  $\|x\| = 1$ , we have thus shown that

$T = M + iN$  is not normal and  $W\{i(MN-NM)\} \geq 0$

(ii) Suppose  $m < 0$ .

For each  $\beta$ , choose

$$c_{\alpha\beta} = \begin{cases} 0 & , \quad \text{if } \alpha = \beta \\ 0 & , \quad \text{if } \alpha \neq \beta , \quad \alpha \geq 3, \quad \beta \geq 3 \end{cases}$$

$$c_{12} > 0 , \quad c_{21} > 0 .$$

With this choice we have  $MN = NM$  so that

$$W\{i(MN-NM)\} = 0 .$$



We shall show that  $T = M + iN$  is not normal because  $M^2$  is not hermitian.

$$M = \begin{pmatrix} ic_{21} & c_{12} \\ c_{21} & ic_{12} \end{pmatrix}$$

Hence

$$M^2 = \begin{pmatrix} c_{21}(c_{12}-c_{21}) & ic_{12}(c_{21}+c_{12}) \\ ic_{21}(c_{21}+c_{12}) & c_{12}(c_{21}-c_{12}) \end{pmatrix}$$

Hence



$$\|I+itM^2\| = \max \begin{cases} |1+itc_{21}(c_{12}-c_{21})| + tc_{21}(c_{21}+c_{12}) , \\ |1+itc_{12}(c_{21}-c_{12})| + tc_{12}(c_{21}+c_{12}) , \\ 1 , \end{cases}$$

$$= \max \begin{cases} 1+o(t)+tc_{21}(c_{21}+c_{12}) , & \text{as } t \rightarrow 0 \\ 1+o(t)+tc_{12}(c_{21}+c_{12}) , & \text{as } t \rightarrow 0 \\ 1+o(t) , & \text{as } t \rightarrow 0 \end{cases} .$$

Since  $c_{21} > 0$  ,  $c_{12} > 0$  ,

$$\|I+itM^2\| \neq 1+o(t) \quad \text{as } t \rightarrow 0 .$$

Hence  $M^2$  is not hermitian.

Thus by a suitable choice of the  $c_{\alpha\beta}$  , we can ensure that

$T = M + iN$  is not normal and  $W\{i(MN-NM)\} \geq 0$  .

Now

$$T = M + iN$$

$$= \begin{pmatrix} (1+i)c_{11} & \\ & (1+i)c_{22} \end{pmatrix} ,$$

$$(1+i)c_{22} ,$$



Suppose  $(S_1, S_2)$  is a pair of subspaces of  $X$  which reduces  $T$  completely.

Then  $X = S_1 \oplus S_2$ .

We may assume without loss of generality that  $S_1$  is spanned by  $\delta^1, \delta^2, \dots, \delta^k$ . Then

$$T|_{S_1} = \begin{pmatrix} (1+i)c_{11} & & & \\ & (1+i)c_{22} & & \\ & & \ddots & \\ & & & (1+i)c_{kk} \end{pmatrix}$$

Thus if  $\lambda = 0$  or  $\lambda \in \sigma(T)$ ,

$$\begin{aligned} \|(T-\lambda)|_{S_1}\| &= \max_{1 \leq \alpha \leq k} \{ |(1+i)c_{\alpha\alpha} - \lambda| \} \\ &= \max [ |\lambda| : \lambda \in \text{P}\sigma\{(T-\lambda)|_{S_1}\} ] \\ &= \max [ |\lambda| : \lambda \in \sigma\{(T-\lambda)|_{S_1}\} ] \\ &= r\{(T-\lambda)|_{S_1}\} . \end{aligned}$$





Similarly we can show that for  $\lambda = 0$  or  $\lambda \in \sigma(T)$  ,

$$\|(T-\lambda)|_{s_2}\| = r\{(T-\lambda)|_{s_2}\} \quad .$$

This shows that  $T = M + iN$  is hyponormal but not normal.



BIBLIOGRAPHY

1. G. Bachman, "Elements of Abstract Harmonic Analysis," Academic Press.
2. E. Berkson, "A characterisation of scalar type operators on reflexive Banach spaces", Pacific Journal of Mathematics, 13 (1963), p. 365-373.
3. E. Berkson, "Some characterisations of  $C^*$ -Algebras," Illinois Journal of Mathematics, Vol. 10(1966), p. 1-8.
4. N. Dunford, "Spectral Theory II. Resolutions of the identity," Pacific Journal of Mathematics, 2(1952), p. 559-614.
5. N. Dunford, "Spectral Operators," Pacific Journal of Mathematics, 4(1954) p. 321-354.
6. N. Dunford, "A survey of the theory of spectral operators," Bull. Amer. Math. Soc., vol. 64(1958), p. 217-274.
7. S. R. Foguel, "The relations between a spectral operator and its scalar part," Pacific Journal of Mathematics, 8(1958), p. 51-65.
8. B. Fuglede, "A commutativity theorem for normal operators," Proc. Nat. Acad., U.S.A., vol. 36(1950), p. 35.
9. B. W. Glickfield, "A metric characterisation of  $C(X)$  and its generalisation to  $C^*$ -Algebras," Illinois Journal of Mathematics, vol. 10(1966), p. 547-556.
10. S. Hildebrandt, "The closure of the numerical range of an operator as a spectral set," Communications in Pure and Applied Mathematics, vol. 17(1964), p. 415-421.
11. E. Hille and R. S. Phillips, "Functional Analysis and Semi-Groups," Amer. Math. Soc. Coll. Publications, vol. 31(1957).
12. G. Lumer, "Semi-inner product spaces," Trans. Amer. Math. Soc., 100(1961), p. 29-43.
13. G. Lumer, "Spectral Operators, hermitian operators and bounded groups," Acta. Sci. Math., Szeged, T. 25(1964), p. 75-85.
14. C. A. McCarthy, "Commuting Boolean algebras of projections," Pacific Journal of Mathematics, 11(1961), p. 295-307.



15. C. E. Rickart, "The General Theory of Banach Algebras," D. Van Nostrand Co., Inc. (1960).
16. F. Riesz and B. Sz. Nagy, "Functional Analysis," F. Ungar Publishing Co. (1955).
17. J. G. Stampfli, "Hyponormal operators," Pacific Journal of Mathematics, 12(1962), p. 1453-1458.
18. A. E. Taylor, "Introduction to Functional Analysis," John Wiley.
19. I. Vidav, "Eine metrische Kennzeichnung der selbstadjungierten operatoren," Math. Zeitschrift, 66(1956), p. 121-128.
20. J. P. Williams, "Spectra of products and numerical ranges," Journal of Math. Anal. and Appl., 17(1967), p. 214-220.





**B29884**